Radiological Case Report / Caso Clínico

Male Breast Hemangioma: A Rare Entity

Hemangioma da Mama no Homem: Uma Entidade Rara

Ricardo Pereira Dias¹, André Peixoto¹, Tiago Oliveira², Dolores Lopez Presa², Inês Leite¹, Leonor Fernandes¹, Isabel Duarte¹, Olga Fouto¹

¹Serviço de Imagiologia Geral, Centro Hospitalar Universitário Lisboa Norte, Lisboa, Portugal

²Serviço de Anatomia Patológica, Centro Hospitalar Universitário Lisboa Norte, Lisboa, Portugal

Address

Ricardo Pereira Dias Serviço de Imagiologia Geral Centro Hospitalar Universitário Lisboa Norte Av. Prof. Egas Moniz 1649-028 Lisboa, Portugal e-mail: pereira.dias.ricardo@gmail.com

Received: 03/05/2020 Accepted: 08/05/2021 Published: 10/09/2021 © Author(s) (or their employer(s)) and ARP 2021. Re-use permitted under CC BY-NC. No commercial re-use.

Abstract

The male breast can be affected by a variety of benign and malignant entities. Breast hemangioma is a rare benign vascular tumor occurring in both genders, being even rarer in men, with only a few cases reported in the literature. This is a case report of a 78-yearold male patient with a known history of advanced laryngeal cancer previously treated with radiotherapy who was admitted to our emergency department in the setting of severe dyspnea. A neck and chest computerized tomography were performed, revealing post radiation changes of the neck and no significant pulmonary abnormalities. An incidental left breast nodule was noted and considered indeterminate. The mammogram and the breast ultrasound revealed a well-circumscribed hyperdense and hypoechoic oval nodule, without spiculations nor calcifications. It also demonstrated a slightly heterogeneous echo pattern with no posterior acoustic shadowing. An ultrasound-guided core needle biopsy was performed and the histopathologic examination was compatible with a cavernous hemangioma.

Keywords

Hemangioma; Breast; Male.

Resumo

A mama masculina pode ser afetada por uma variedade de entidades benignas e malignas. O hemangioma da mama é um tumor vascular benigno raro que pode ocorrer em ambos os sexos, sendo ainda mais raro em homens, com poucos casos relatados na literatura. Apresenta-se um caso de um homem de 78 anos com antecedentes de cancro da laringe avançado, previamente tratado com radioterapia, que recorreu ao nosso serviço de urgência por dispneia grave. Realizouse uma tomografia computadorizada do pescoço e do tórax que revelou alterações pósradicas cervicais, sem alterações pulmonares significativas. Um nódulo incidental na mama esquerda foi também revelado e considerado indeterminado. A mamografia e ecografia realizadas demonstraram um nódulo oval, hiperdenso e hipoecogénico, bem circunscrito, sem espiculações ou calcificações, demonstrando uma ecoestrutura ligeiramente heterogénea, sem sombra acústica posterior. Foi realizada microbiópsia ecoguiada e o exame histopatológico foi compatível com um hemangioma cavernoso, de tipo sinusoidal.

Palavras-chave

Hemangioma; Mama; Homem.

Case presentation

A 78-year-old male with a medical history of laryngeal squamous cell carcinoma, cT4aN1M0 stage treated with radiotherapy, was referred to the emergency department of our institution due to exacerbated dyspnea, prompting the placement of a tracheostomy. In this setting, he underwent a contrast-enhanced computerized tomography (CT) which revealed diffuse soft-tissue post radiotherapy edema of the neck resulting in upper airway obstruction. Neither signs of locoregional recurrence nor significant pulmonary abnormalities were noted. However, a slightly enhancing oval and well circumscribed soft-tissue nodule was noted on the left breast measuring 25 mm (Fig. 1). It was considered indeterminate and further breast imaging evaluation was recommended.

The patient had a history of smoking and alcohol consumption. Besides bilateral breast enlargement due to gynecomastia, the patient had no breast pain nor nipple discharge. He denied family history of breast or ovarian cancer. On clinical examination, a small, soft and slightly mobile nodule was palpable in the upper outer quadrant of the left breast. The nodule was not attached to the chest wall. The mammogram revealed a well-defined hyperdense nodule in the left upper outer quadrant without microcalcifications or spiculations along with bilateral gynecomastia (Fig. 2A). The lesion appeared on ultrasound as an oval hypoechoic and slightly heterogeneous lesion with smooth margins and no posterior shadowing measuring 26 mm (Fig. 2B). There was no axillary lymphadenopathy. The imaging findings were consistent with a BI-RADS 4 score and an ultrasound-guided needle-core biopsy was performed. The histopathologic examination of the lesion was compatible with a cavernous hemangioma of the sinusoidal type (Fig. 3). The lesion remained stable after a 12-month follow-up.

Discussion

Hemangiomas are benign vascular tumors that rarely develop in the breast. Perilobular hemangiomas are benign and microscopic lesions without clinical significance, generally smaller than 2 mm (not apparent on mammography).^{1,2,3,4} Breast hemangiomas that are clinically



Fig. 1 – Contrast-enhanced chest CT showing a well-circumscribed, slightly enhancing, and homogeneous mass located in the upper outer quadrant of the left breast (white arrow).



Fig. 2 – Mediolateral oblique views of the mammogram revealing bilateral gynecomastia and a well circumscribed and lobulated dense mass located at the upper outer quadrant of the left breast, without spiculations, calcifications, or skin retraction (A, white arrow). Left breast ultrasound images demonstrating a 26 mm oval solid mass with well-circumscribed margins and parallel orientation with slightly complex echotexture but without posterior acoustic shadowing (B, yellow arrow).



Fig. 3 – Hemangioma (A – hematoxylin-eosin, x100; B/C – CD31/Podoplanin, x100). The lesion was composed of thin-walled cavernous vessels that interconnect in a sinusoidal pattern (A). These vessels are lined by bland endothelial cells with no cytological atypia. The cells showed immunoreactivity for CD31 (B) and negativity for Podoplanin (C).

and/or mammographically detectable are classified into cavernous, capillary, complex or venous types, depending on the size of their vascular channels. The cavernous type hemangioma is the most frequently encountered.

In a previous study comprising 1362 patients of both genders treated with lumpectomy, simple mastectomy or modified radical mastectomy for various primary breasts tumors, only 12 cases of hemangiomas were found (11 women and one man; 0,8% of total breast lesions).⁴ As the number of male patients enrolled in the study population was not specified, the real incidence of male breast hemangiomas remains unclear. Although imaging features of breast hemangiomas are exceptionally rare and only few cases have been previously reported.

Breast hemangiomas tend to be larger in men (mean diameter of 5.4 cm) than in women (mean diameter of 1.1 cm).⁵ Apart from size, the other radiologic features of male breast hemangiomas show no differences from those reported in women.^{5,6,7,8} Breast hemangiomas usually appear as superficial, oval and well circumscribed nodules or masses and, in some cases, can be slightly lobulated. They commonly have a density similar to that of fibroglandular breast tissue in mammograms and lack spiculations.^{5,6,7,8} Although uncommon, cases presenting with fine or coarse calcifications have been reported.9 Variable echo patterns are seen on ultrasound. Around one-third of hemangiomas exhibit hyperechoic echotexture and the remaining may show an isoechoic, hypoechoic or a complex echo pattern. A common feature is the absence of a posterior acoustic shadow.67,10 Vascular poles may be seen with Doppler ultrasound.¹⁰ Magnetic Resonance Imaging (MRI) usually demonstrates a lesion isointense to the surrounding fibroglandular tissue on T1-weighted images and hyperintense on T2-weighted images.7 Peripheral arterial enhancement and delayed central enhancement may be seen on contrast-enhanced images - a useful imaging feature that may aid in differentiating it from malignant tumors.7

The differential diagnosis of breast hemangiomas in men is broad but should include angiosarcoma, metastasis (particularly in patients with a known primary cancer), breast cancer and pseudoangiomatous stromal hyperplasia (PASH).^{9,10,11,12,13}

Angiosarcoma, although extremely rare, is considered the main differential diagnosis since this vascular malignant tumor frequently mimics the imaging appearance of a breast hemangioma on mammography and ultrasound.6 MRI may provide some helpful clues in some cases. Indeed, high-grade angiosarcomas tend to present rapid enhancement and washout with frequent visualization of large draining vessels and multiple unenhanced areas within the tumor.¹¹ However, histopathologic analysis remains mandatory for correct diagnosis. Microscopically, both can be composed of thin-walled capillaries lined by low nuclear grade endothelial cells and both are typically positive for vascular markers such as CD31 and CD34 but negative for lymphatic endothelium marker, podoplanin.8 However, angiosarcomas are positive for the S-phase kinase-associated protein 2, show a higher Ki-67 proliferation index and other

features of architectural and/or cytological atypia normally absent in breast hemangiomas, as well as a frequent previous history of chest irradiation, usually in the context of breast cancer treatment. In contrast to angiosarcomas, hemangiomas also usually present well-circumscribed vascular channels surrounding the ducts and lobules rather than invading them.⁸ Currently, it is considered that with clinical, radiologic and pathologic correlation, core needle biopsies allow for a reliable histopathological diagnosis. Thus, further surgical excision is no longer mandatory and a follow-up may be preferred, especially in the case of frail patients that would not be good candidates for surgery,¹¹ such as ours.

Metastases to the breast are extremely rare, representing 0.5%-3% of all malignant breast tumors concerning both genders.8 The most common primary malignancies that metastasize to the breast are melanoma, non-Hodgkin's lymphoma, and carcinomas of the lung, stomach, kidneys, prostate and ovaries.8 Breast metastasis from squamous cell carcinoma from the head and neck is highly uncommon, with only a few cases reported.14 Hematogeneous breast metastases are frequently multiple or bilateral round masses with circumscribed or microlobulated margins and no associated calcifications on mammography.8,13 Nevertheless, single irregular masses with spiculated margins and calcifications may also occur.15 Posterior acoustic enhancement is a common finding on ultrasound, with variable vascularity on Colour Doppler.¹⁵ Metastatic lesions are typically isointense on MRI in both T1 and T2-weighted images, with homogeneous enhancement on post-contrast images.15 Lymphatic spread presents as diffuse oedema with skin and trabecular thickening and is commonly accompanied by pathologic axillary lymph nodes.^{8,13,15}

Male breast cancer is rare, accounting for <1% of male cancers and <1% of all breast cancers13. Invasive breast carcinoma of no special type (NST) is the most frequent histological type.^{8,13} As no cancer screening program exists for men, male breast cancer is frequently diagnosed at an advanced stage.13 Contrary to breast hemangiomas, breast male cancers usually present with suspicious imaging features such as irregular, spiculated or indistinct margins on both mammography and ultrasound, and they are frequently located in the subareolar region or eccentric to the nipple.^{8,13} Contrary to women, the concept of benign calcifications does not apply to a male breast mass and should always be considered suspicious for breast cancer.14 PASH is a frequent benign entity (found in 50 % of male breast biopsies) consisting of a hormonally-dependent proliferation of fibroblasts and myofibroblasts, being highly associated with gynecomastia.8,13 Although variable, the most common imaging finding is a non-calcified oval, circumscribed mass at mammography and an ovoid homogenously hypoechoic mass at ultrasound.^{8,13}

Hemangiomas are extremely rare and benign in male breasts but share similar imaging features with those appearing in women. While ultrasound and MRI may provide useful clues that aid in the differential diagnosis, a histopathological examination remains mandatory for proper diagnosis and management.

Ethical disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare. Conflitos de interesse: Os autores declaram não possuir conflitos de interesse. Financing Support: This work has not received any contribution, grant or scholarship.

Suporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients. *Confidencialidade dos dados:* Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Jozefczyk MA, Rosen PP. Vascular tumors of the breast. II. Perilobular hemangiomas and hemangiomas. Am J Surg Pathol. 1985;9:491-503.

2. Rosen PP, Hoda SA, Brogi E, Koerner FC. Rosen's breast pathology. Fourth edition. Philadelphia: Wolters Kluwer Health - Lippincott Williams & Wilkins, 2014.

 Lesueur GC, Brown RW, Bhathal PS. Incidence of perilobular hemangioma in the female breast. Arch Pathol Lab Med. 1983;107:308-10.
Kondi-Pafiti A, Dellaportas D, Myoteri D, Tsagkas A, Ntakomyti E, Kairi-Vasilatou E. Rare non-epithelial primary breast neoplasms: a tenyear experience at a Greek University Hospital. J BUON. 2013;18:70-6.

5. Yoga A, Lyapichev KA, Baek D, Gomez-Fernandez CR. Hemangioma of a male breast: case report and review of the literature. Am J Case Rep. 2018;19:1425-9.

6. Mesurolle B, Sygal V, Lalonde L, Lisbona A, Dufresne MP, Gagnon JH, et al. Sonographic and mammographic appearances of breast hemangioma. Am J Roentgenol. 2008;191:W17-22.

 Glazebrook KN, Morton MJ, Reynolds C: Vascular tumors of the breast: mammographic, sonographic, and MRI appearances. Am J Roentgenol. 2005;184:331-8.

8. Lattin GE Jr, Jesinger RA, Mattu R, Glassman LM. From the radiologic pathology archives: diseases of the male breast: radiologic-pathologic correlation. Radiographics. 2013;33:461-89.

9. Shi AA, Georgian-Smith D, Cornell LD, Rafferty EA, Staffa M, Hughes K, et al. Radiological reasoning: male breast mass with calcifications. Am J Roentgenol. 2005;185:S205-10.

10. Isaza Zapata S, Uribe González R, González Vasquez CM. Breast hemangioma in a male patient: report of a case. Radiol Case Rep. 2019;14:427-9.

11. Wu WH, Ji QL, Li ZZ, Wang QN, Liu SY, Yu JF. Mammography and MRI manifestations of breast angiosarcoma. BMC Womens Health. 2019;19:73.

12. Mango VL, Goodman S, Clarkin K, Wynn RT, Friedlander L, Hibshoosh H, et al. The unusual suspects: a review of unusual benign and malignant male breast imaging cases. Clin Imaging. 2018;50:78-85.

13. Chesebro AL, Rives AF, Shaffer K. Male Breast Disease: What the Radiologist Needs to Know. Curr Probl Diagn Radiol. 2019;48:482-93.

14. Longo R, Melgar E, Campitiello M, Plastino F, Eid N, Quirin I, et al. Breast metastasis from squamous cell carcinoma of the oropharynx: a case report. J Med Case Rep. 2017;11:355.

15. Sippo DA, Kulkarni K, Carlo PD, Lee B, Eisner D2, Cimino-Mathews A, et al. Metastatic disease to the breast from extramammary malignancies: a multimodality pictorial review. Curr Probl Diagn Radiol. 2016;45:225-32.