

Radiological Case Report / Caso Clínico

Prophylactic Balloon-Occlusion of the Hypogastric Arteries in Surgical Treatment Placenta Accreta Spectrum Disorders: How We Do It

Oclusão com Balão das Artérias Hipogástricas no Tratamento Cirúrgico do Acretismo Placentário: A Nossa Abordagem

Raquel Gaio¹, Maria Pulido Valente², Maria de Carvalho Afonso², Ana Isabel Simões Ferreira¹

¹Department of Radiology, Hospital de Santa Maria, Centro Hospitalar Universitário de Lisboa Norte, E.P.E., Lisboa, Portugal
²Department of Obstetrics, Gynecology and Reproductive Medicine, Hospital de Santa Maria, Centro Hospitalar Universitário Lisboa Norte, E.P.E., Lisboa, Portugal

Address

Raquel Coutinho de Sousa Gaio
Rua Silva e Albuquerque 17, 4º
1700-360, Lisboa, Portugal
e-mail: gaio.sousa.raquel@gmail.com

Received: 11/05/2021

Accepted: 08/07/2021

Published: 10/09/2021

© Author(s) (or their employer(s)) and ARP 2021. Re-use permitted under CC BY-NC. No commercial re-use.

Abstract

Placenta accreta spectrum disorders are amongst the most dangerous conditions associated with pregnancy which can lead to massive obstetric hemorrhage and, secondarily, to coagulopathy, multisystem organ failure and maternal death. Perioperative endovascular balloon occlusion of the pelvic vessels has been shown to be effective in reducing estimated blood loss, transfusion requirements, and in providing favorable surgical conditions, with low procedure-related complications. Therefore, the interventional radiologist should be familiar with the indications and technical considerations of the procedure. The optimal positioning for occlusion balloon catheters is still controversial and protocols differ between institutions and local expertise. Using a case of placenta increta as example, this report provides detailed description of the technical details and materials used in our center for preoperative prophylactic balloon-occlusion of the hypogastric arteries (our preferred approach), from the angio suite until completion of hysterectomy.

Keywords

Placenta accreta; Placenta diseases; Obstetric labor complications; Balloon occlusion; Interventional radiology.

Resumo

A morbimortalidade associada ao acretismo placentário resulta da hemorragia obstétrica massiva e das suas complicações secundárias, como coagulopatia, falência multiorgânica e morte materna. A oclusão peri-cirúrgica dos vasos pélvicos com balões endovasculares mostrou ser eficaz na redução das perdas hemáticas, necessidade transfusional e melhoria das condições cirúrgicas, com baixa incidência de complicações associadas ao procedimento. Assim, o radiologista de intervenção deve estar familiarizado com as indicações e considerações técnicas deste procedimento. O posicionamento ideal dos balões oclusivos é, todavia, ainda controverso e os protocolos diferem entre as instituições e a experiência local. Usando um caso de placenta increta como exemplo, este artigo fornece uma exposição detalhada dos pormenores técnicos e materiais utilizados no nosso centro para a oclusão com balão das artérias hipogástricas (a nossa abordagem preferencial), desde a sala de angiografia até à conclusão da histerectomia no bloco operatório.

Palavras-chave

Placenta accreta; Doenças da placenta; Complicações obstétricas do parto; Radiologia de intervenção.

Introduction

Morbimortality associated with placenta accreta spectrum (PAS) disorders occur mostly during childbirth when trying to remove the placenta from the uterus, which can trigger massive obstetric hemorrhage.¹

Timely delivery planning and management by an experienced multidisciplinary team have been shown to improve maternal and neonatal outcomes.² In tertiary-care centers, like ours, minimally invasive endovascular interventional radiology (IR) techniques performed before surgery have become an integral part of management on women affected by PAS disorders.^{3,4} They have been shown to be effective in reducing estimated blood loss, transfusion requirements, and in providing favorable surgical conditions for the gynecology and obstetrics (GO) team (better hemostasis and field visibility). Literature data is, however, heterogeneous and most studies have poor quality.^{4,5} The optimal positioning for occlusion balloon (OB) catheters is also controversial and protocols differ

between institutions and local expertise. Our preference is for perioperative prophylactic balloon-occlusion of the hypogastric arteries (PBOHA).

The procedure was first done in our institution in 2015 and rarely performed until 2018. Since then, the GO and IR teams started working together in developing a protocol to optimize the response to these patients. A total of 10 cases with PAS disorder were submitted to PBOHA before planned cesarean delivery and all performed hysterectomy, except one. Although the protocol is at an initial stage, so far, outcomes have been positive, within literature references and no endovascular complications were registered.

Aiming to share our favorable experience, this report provides a detailed description of the technical aspects and materials used in our center for PBOHA, from the angio suite until completion of hysterectomy. For that, we use a case of a pathologically proven placenta increta as an example.

Case Report

A 42-year-old woman with placenta previa was admitted to our hospital at 25 weeks of gestation due to vaginal bleeding. This was the third ongoing pregnancy with two previous cesarean deliveries. Ultrasound and magnetic resonance imaging, at 28 and 30 weeks of gestation, respectively, showed complete placenta previa and positive findings for invasive placentation.

Surgical and perioperative multidisciplinary planning was done by GO, anesthesiology, IR, neonatology and immunohemotherapy teams. An elective cesarean was scheduled at 36 weeks with possible need for total hysterectomy. Due to the high risk of surgical bleeding, additional PBOHA was planned.

On the scheduled day, the patient went first to the angio suite for placement of OB in the hypogastric arteries (HA). Patient sedation and analgesia was managed by the anesthesiologists' team. After local anesthesia (2% lidocaine), bilateral common femoral artery access was obtained, using 6F sheaths (*Radifocus® Introducer II Guidewire-J Spring, 6 F, Terumo*). Manual injection of contrast through one of the sheaths was used to delineate both HA anatomy and obtain a road map. An hydrophilic Cobra 2 catheter (*Glidecath®, 5 F, 100 cm, Terumo*) and an hydrophilic angled tip guidewire (*Glidewire® Hydrophilic Coated Guidewire, 0,035", 180 cm, Terumo*) were used to select the contralateral HA. An OB (*Advance® 7x20mm, 0.035", 5 F, 80 cm, Biosonda-Cook*) was then positioned over-the-wire, in the proximal portion of the HA, under intermittent fluoroscopic view; the road map was only used to confirm the final position of the balloon. After satisfactory positioning, the OB was tested by inflating, until nominal pressure was achieved, and, afterwards, deflated. The procedure was then repeated for the contralateral side. A final image was obtained with bone reference, to allow for balloon position confirmation in the operating room (Fig. 1). Total fluoroscopy time was 3.9 minutes, dose area product (DAP) was 28.215 Gy*cm² and air kerma was 104 mGy. Lastly, guidewires were secured to the OB catheters by unidirectional hemostatic valves (*Flo30™, MeritMedical*), sheaths were sutured to the

skin, and both sheaths and OB catheters were securely fixed together using a *Skater™Fix (Argon)*; the whole material was then wrapped in two separate sterile fields before the patient was transferred to the operating room (Fig. 2).

Once the patient arrived at the operating room, a big sterile field was positioned under the IR material. The OB catheters were straightened along patients' legs and attached to the inflation pumps, allowing the interventional radiologist to have rapid access from the bottom, without disturbing surgeons. Fluoroscopic confirmation of the OB position was done. The patient was then routinely prepared for cesarean section. At surgery, a bulging placenta in the anterior uterine wall was confirmed, although there were no signs of bladder invasion. Uterine incision was performed above the upper limit of the bulging placenta to avoid it. After fetal extraction, both OB were inflated to their nominal pressure. A gentle attempt to remove the placenta was unsuccessful, so it was left in situ. Hysterectomy with bilateral salpingectomy was performed and OB were inflated for 40 minutes. During deflation no active bleeding was seen by the surgeons. After surgical incision was closed, OB catheters were removed under fluoroscopic control and puncture sites were occluded with a vascular closure device (*Exoseal®, 6 F, Cordis*).

During surgery, tranexamic acid and one pack of red blood cells were given. The total estimated blood loss was 900cc. The lowest value of hemoglobin was 8,9g/dL and no further blood transfusions were needed. The patient was transferred to the intensive care unit for hemodynamic monitoring in the early postoperative period. No complications were reported. The newborn boy was healthy, weighed 2540g and the APGAR score was 9 and 10 at the first and fifth minutes, respectively. The patient was discharged at the sixth day postpartum. Anatomopathological examination of the uterus showed loss of the decidual layer and deep invasion of the myometrium, less than 1 mm away of the serosa compatible with placenta increta.




Figure 1 – Fluoroscopic images at the angio suite. A) Initial angiogram to delineate HA anatomy and road mapping guidance. B and C) Test inflation of the left occlusion balloon using diluted water-soluble contrast. Note that it is not possible to inject contrast to control the second balloon. D) Final image with bone reference, to guide balloon inflation in the operating room. Note the position of the guidewire in branches of the posterior division of the hypogastric artery.




Figure 2 – Preparation of the vascular access sites. The sheaths of the balloons were secured to the skin using a *Skater™Fix* (asterisks in A) and the whole material was then wrapped in two separate sterile dressings (white arrows in B).

Discussion

Planned cesarean hysterectomy, leaving the placenta in situ, is the general recommendation from the American College of Obstetricians and Gynecologists and remains the definitive surgical treatment for PAS disorders, especially for its invasive forms.^{1,5} Minimally invasive IR techniques allow for preoperative placement of endovascular balloons to occlude the bloodstream proximally to the uterine arteries and promote a decrease in uterine arterial supply, after umbilical cord clamping. The main goal is to reduce hemorrhage; even when successful vessel occlusion is achieved, some blood loss is expected to occur.⁵

The endovascular approach in the angio suite is not long and has a low rate of adverse events;^{4,7} nevertheless, the IR team has to be available to accompany the patient through the whole process. The placement of the OB is done by the two most experienced interventional radiologists, as well as a trained technician operating the angiography equipment, to minimize possible side effects. There is concern to minimize radiation to the fetus, using intermittent pulse and stored fluoroscopy (instead of acquisitions), with collimated images, no magnification and low pulse rates.^{8,9,10} The optimal position of OB catheters placement is still controversial and heterogeneous results are found when reviewing the latest literature. Arterial occlusion upstream of the HA (at the common iliac arteries or infrarenal aorta) has the potential advantage of blocking vascular flow at the collateral pathways between the hypogastric and external iliac arteries. This is however counterbalanced by the potential risk of greater traumatic injury and ischemic damage of a larger vascular territory.^{5,10} Although HA balloon occlusion does not prevent collateral flow from external iliac branch, the vascular territory at risk of perfusion injury is smaller and trauma is diminished by the use of smaller-caliber material. Though opinions and protocols may differ between institutions and local expertise, the latest systematic review and meta-analysis concerning this topic wasn't able to demonstrate a meaningful comparison between different IR techniques, mainly due to the small sample size of most studies.⁴ Our preference is for placement of the OB bilaterally in the proximal segment of the HA by contralateral femoral approach.

The size of the balloon should be optimally tailored to the vessel being occluded. The MRI done preoperatively

is a useful tool to predict the size of the OB that will be needed in advance. We usually upgrade the diameter of the OB by 1-2 mm of the vessel caliber, bearing in mind that after delivery the size of the artery tends to increase slightly. Careful must be taken to never inflate both OB at the same time before delivery, as this would considerably decrease blood flow for the fetus. After childbirth, if necessary (when blood loss is higher than expected) we may additionally inflate the OB 1-2 mmHg above nominal pressure, to compensate for a slight increase of the artery caliber that can occur after delivery. As illustrated in this case, we routinely use 20mm *Advance*® 0.035", *Biosonda-Cook* OB catheters of 7 mm, 8 mm or 9 mm of diameter, and we have been able to achieve maximal wall tension without vessel rupture.

During OB placement, the tips of the guidewires should preferentially be kept in the posterior division of the HA to avoid potential spasm of the anterior division and its uterine branch. The guidewires are kept inside the OB catheters to confer more stability to the latter and to allow for rapid repositioning of the OB in case they are dislodged.

Some centers leave the OB inflated after surgery, bearing in mind the increased risk of hemorrhagic complications in the following hours.^{5,11} Similarly to other centers,^{9,12,13,14} we deflate the OB after surgical procedure is finished OB (maximum OB inflation time in our experience was 60 minutes). Even though there is, once more, no consensus on this matter, our approach relates to safety concerns (lack of staff to tightly monitor possible complications). Thrombosis and reperfusion injury of the lower limbs, although uncommon, remain the main complications of the endovascular procedure).^{5,11} Postoperative care includes absolute rest for 4 hours, avoidance of physical efforts for 48 hours, inguinal bandage surveillance for the first 12 hours and vital signs monitoring.

In conclusion, PBOHA should be considered as part of a multidisciplinary approach to reduce postpartum uterine hemorrhage in the setting of PAS disorders. It is a minimally invasive endovascular procedure, uncommonly associated with adverse events in experienced hands. It has been easily implemented in our center, and although it requires full availability from the IR team, it does not add a great deal of complexity to the surgical approach.

Ethical disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare.

Conflitos de interesse: Os autores declaram não possuir conflitos de interesse.

Financing Support: This work has not received any contribution, grant or scholarship.

Suporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Confidencialidade dos dados: Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Practice ACoO. ACOG Committee Opinion. Placenta accreta. No. 266, January 2002. American College of Obstetricians and Gynecologists. *Int J Gynaecol Obstet.* 2002;77:77-8.
2. Shamshirsaz AA, Fox KA, Erfani H, Clark SL, Salmanian B, Baker BW, Coburn M, Shamshirsaz AA, Bateni ZH, Espinoza J, Nassr AA, Popek EJ, Hui SK, Teruya J, Tung CS, Jones JA, Rac M, Dildy GA, Belfort MA. Multidisciplinary team learning in the management of the morbidly adherent placenta: outcome improvements over time. *Am J Obstet Gynecol.* 2017;216:612.e1-612.e5.
3. Shahin Y, Pang CL. Endovascular interventional modalities for haemorrhage control in abnormal placental implantation deliveries: a systematic review and meta-analysis. *Eur Radiol.* 2018;28:2713-26.
4. D'Antonio F, Iacovelli A, Liberati M, Leombroni M, Murgano D, Cali G, Khalil A, Flacco ME, Scutiero G, Iannone P, Scambia G, Manzoli L, Greco P. Role of interventional radiology in pregnancy complicated by placenta accreta spectrum disorder: systematic review and meta-analysis. *Ultrasound Obstet Gynecol.* 2019;53:743-51.
5. Angileri SA, Mailli L, Raspanti C, Ierardi AM, Carrafello G, Belli AM. Prophylactic occlusion balloon placement in internal iliac arteries for

- the prevention of postpartum haemorrhage due to morbidly adherent placenta: short term outcomes. *Radiol Med.* 2017;122:798-806.
6. Allen L, Jauniaux E, Hobson S, Papillon-Smith J, Belfort MA; FIGO Placenta Accreta Diagnosis and Management Expert Consensus Panel. FIGO consensus guidelines on placenta accreta spectrum disorders: Nonconservative surgical management. *Int J Gynaecol Obstet.* 2018;140:281-90.
7. Shahin Y, Pang CL. Endovascular interventional modalities for haemorrhage control in abnormal placental implantation deliveries: a systematic review and meta-analysis. *Eur Radiol.* 2018;28:2713-26.
8. Dauer LT, Thornton RH, Miller DL, Damilakis J, Dixon RG, Marx MV, Schueler BA, Vañó E, Venkatesan A, Bartal G, Tsetis D, Cardella JF. Society of Interventional Radiology Safety and Health Committee; Cardiovascular and Interventional Radiology Society of Europe Standards of Practice Committee. Radiation management for interventions using fluoroscopic or computed tomographic guidance during pregnancy: a joint guideline of the Society of Interventional Radiology and the Cardiovascular and Interventional Radiological Society of Europe with Endorsement by the Canadian Interventional Radiology Association. *J Vasc Interv Radiol.* 2012;23:19-32.
9. Carnevale FC, Kondo MM, de Oliveira Sousa W Jr, Santos AB, da Motta Leal Filho JM, Moreira AM, Baroni RH, Francisco RP, Zugaib M. Perioperative temporary occlusion of the internal iliac arteries as prophylaxis in cesarean section at risk of hemorrhage in placenta accreta. *Cardiovasc Intervent Radiol.* 2011;34:758-64.
10. Nankali A, Salari N, Kazemina M, Mohammadi M, Rasoulinya S, Hosseinian-Far M. The effect of prophylactic internal iliac artery balloon occlusion in patients with placenta previa or placental accrete spectrum: a systematic review and meta-analysis. *Reprod Biol Endocrinol.* 2021;19:40.
11. Teixidor Viñas M, Chandrarahan E, Moneta MV, Belli AM. The role of interventional radiology in reducing haemorrhage and hysterectomy following caesarean section for morbidly adherent placenta. *Clin Radiol.* 2014;69:e345-51.
12. Kaufman C, Tadros A. Endovascular Interventions for the Morbidly Adherent Placenta. *J Clin Med.* 2018;7:92.
13. Picel AC, Wolford B, Cochran RL, Ramos GA, Roberts AC. Prophylactic Internal Iliac Artery Occlusion Balloon Placement to Reduce Operative Blood Loss in Patients with Invasive Placenta. *J Vasc Interv Radiol.* 2018;29:219-224.
14. Petrov DA, Karlberg B, Singh K, Hartman M, Mittal PK. Perioperative Internal Iliac Artery Balloon Occlusion, In the Setting of Placenta Accreta and Its Variants: The Role of the Interventional Radiologist. *Curr Probl Diagn Radiol.* 2018;47:445-451.