

Radiological Case Report / Caso Clínico

Pelvic Actinomycosis

*Actinomicose Pélvica*José Veiga¹, Margarida Varela dos Santos², João Lopes Dias^{1,3}¹Serviço de Radiologia, Centro Hospitalar Universitário Lisboa Central, Lisboa, Portugal²Serviço de Anatomia Patológica, Centro Hospitalar Universitário Lisboa Central, Lisboa, Portugal³Serviço de Radiologia, CUF Tejo, Lisboa, Portugal

Address

José Veiga
 Serviço de Radiologia
 Centro Hospitalar Universitário Lisboa Central, Hospital Santa Marta
 Rua de Santa Marta 50
 1169-024, Lisboa, Portugal
 e-mail: jveiga92@gmail.com

Received: 29/07/2021

Accepted: 17/04/2022

Published: 31/08/2022

© Author(s) (or their employer(s)) and ARP 2022. Re-use permitted under CC BY-NC. No commercial re-use.

Abstract

Actinomycosis is a rare chronic infection, caused by gram-positive bacteria from the *Actinomyces* species, which is part of the endogenous flora of the oral cavity, gastrointestinal and female genital tracts. In order to cause infection, disruption of the mucosal barrier is imperative. Long-standing IUD (intra-uterine device) is a risk factor. We, hereby, present a case of a 32-year-old woman who presented with lower abdominal pain and a tender mass. The inflammatory parameters increased progressively. There was a history of IUD usage. Ultrasound, Computed Tomography and Magnetic Resonance Imaging demonstrated a heterogeneous enhancing mass-like extending from the bladder to the rectus abdominis muscle, with restricted diffusion and extensive regional oedema. Aspiration cytology revealed bacteria *Parvimonas micra* and polymorphonuclear leukocytes. However, the patient underwent surgery and the histopathological findings showed an abscess with microorganisms compatible with *Actinomyces*.

Key-words

Actinomycosis; Actinomyces; Bacterial infection; Pelvic infection; Intrauterine Device; Magnetic resonance imaging.

Resumo

A actinomicose é uma infeção crónica rara, causada por bactérias gram-positivas da espécie *Actinomyces*, integrantes da flora endógena da cavidade oral, gastrointestinal e genital feminina. Para causar infeção é necessário existir uma disrupção da mucosa. A presença de um dispositivo intra-uterino (DIU) de longa duração é um factor de risco. Apresentamos um caso de uma mulher de 32 anos que recorre ao Serviço de Urgência com dor abdominal, massa hipogástrica dolorosa e aumento progressivo dos parâmetros inflamatórios. Tinha como antecedente de relevo a utilização de um DIU. Foram realizadas ecografia, Tomografia Computadorizada e Ressonância Magnética que demonstraram uma lesão desde a cúpula vesical até ao músculo recto abdominal, com captação heterogénea e restrição à difusão, associada a marcado edema regional. A citologia aspirativa identificou a bactéria *Parvimonas micra* e leucócitos polimorfonucleares. Foi submetida a ressecção cirúrgica com resultado anatomopatológico de abscesso por *Actinomyces*.

Palavras-chave

Actinomicose; Actinomyces; Infeção bacteriana; Infeção Pélvica; Dispositivo intra-uterino; Ressonância Magnética.

Clinical Case

A previously healthy 32-year-old woman was admitted to the emergency room with lower abdominal pain. The pain began 3 days before and was aggravated by abdominal wall muscles contraction. Dysuria was also mentioned. There was no fever.

She was taking oral contraceptives, but she had a history of IUD (intra-uterine device) and IUS (intra-uterine system) rejection.

At physical examination, a tender mass in the hypogastric region/left iliac fossa was palpable.

Laboratory findings only revealed elevated PCR (42,8 mg/L). There was no leucocytosis and urine analysis was unremarkable.

An ultrasound and CT were performed in the emergency room (Figure 1).

The patient underwent MRI in the following week (Figure 2).




Figure 1 – (a) ultrasound image with Power-doppler showing heterogeneous hypoechoic, ill-defined and serpiginous tract, extending from the anterosuperior bladder (blue star) wall to the rectus abdominis muscle (orange dashed lines); (b) – corresponding axial CT image with heterogeneous enhancing mass-like extending from the bladder to the rectus abdominis muscle (orange dashed lines).




Figure 2 – pelvic MRI showing ill-defined mass (red dashed line) in the extraperitoneal cavity measuring 7x5x4 cm, invading the left rectus abdominis muscle and focally the dome of the bladder, with intermediate T2 signal (a) and high T2FS signal compatible with oedema (b), there is also extensive oedematous infiltration of the surrounding fatty tissue from the space of Retzius to the presacral space. The mass shows restricted diffusion (green arrows) - low ADC value (c) and high signal in b1000 (d) – due to either abscess or tumor infiltration. In the post-gadolinium sagittal image (e) the enhancement of the lesion walls and the absent enhancement of the central part is visible. The ovaries (yellow stars) have normal appearance and size, showing multiple physiologic follicles (a).

Inflammatory parameters were increased (RCP - 247 mg/L and leucocytosis - $17 \times 10^9/L$ with neutrophilia 80%). Aspiration cytology revealed gram-positive bacteria (*Parvimonas micra*) and polymorphonuclear leukocytes. The patient underwent surgical resection and the histopathology findings were granulation tissue and an inflammatory process with abscess in the cellular

subcutaneous tissue and trans-mural extension in the bladder, showing microorganisms compatible with *Actinomyces*. (Figure 3) She was discharged with antibiotherapy (doxycycline 100mg q.i.d. during 10 days, then switched to amoxicillin 1g t.i.d. due to gastrointestinal side effects). Follow-up CT scan showed no relevant abnormalities (Figure 4).




Figure 3 (a-b) - (a) A large colony of *Actinomyces* (dark arrowhead) within an abscess, enveloped by an abundant inflammatory infiltrate. Haematoxylin-eosin, 400x. (b) *Actinomyces* colonies can present with a darkly staining eosinophilic rim (dark arrowheads) at the periphery of the colony referred to as the Splendore-Hoeppli phenomenon, which mainly consists of immunoglobulin and cell debris and can occur around colonies of fungi, bacteria and other parasitic organisms. The colony is surrounded by a dense inflammatory infiltrate, in which neutrophils (white arrowheads) are the main component. Haematoxylin-eosin, 600x.



Figure 4 – Post-contrast CT in the sagittal (a) and axial planes (b-c). The images demonstrate the resolution of the previously seen abscess. Residual mild anterior bladder wall thickening is noted (orange arrows).

Discussion

Actinomycosis is a rare chronic infection, caused by gram-positive bacteria from the *Actinomyces* species. *Actinomyces* is part of the endogenous flora of the oral cavity, gastrointestinal tract, and female genital tract. As a result, the frequency of involvement is 50-65% in the cervicofacial area, 15-30% thoracic and 20% abdominopelvic.¹

In order to cause infection, disruption of the mucosal barrier is imperative, which can be caused by trauma, surgery or foreign bodies.²

Long-standing IUD is a risk factor extensively mentioned in the literature,³ however, in our case the patient had one IUD for 6 months and one IUS for 1 month, both rejected. We have not found correlation between actinomycosis and IUD rejection in the literature. We hypothesize that microtrauma occurring during the rejection process is a predisposing factor for infection.

Disease may occur months or years after the removal of an IUD.¹

Pelvic actinomycosis is a difficult radiologic diagnosis, often indistinguishable from malignancy. In our case, the differential diagnosis was malignancy of the urachus, due to its topography.

The disease can spread extensively to the uterus, urinary bladder, rectal area, urachus, abdominal wall and peritoneum. It is also characterised by transfascial spread and sinus tracts.¹

Rim-enhancing abscesses within the mass and extensive inflammatory extensions are usually present. Due to the characteristic presence of fibrosis, solid parts of the mass and the surrounding soft-tissue strands tend to show strong contrast enhancement.⁴

Regional lymphadenopathy and ascites are usually absent, which helps in the distinction from malignancy.⁵

Most actinomycotic infections are polymicrobial in nature,⁶ which justifies the identification of *Parvimonas micra* in the cytology of our patient.

Microorganism diagnosis may be obtained by microscopic examination and the presence of sulphur granules stained with a 1% methylene blue solution.³

Histopathology shows suppurative and granulomatous inflammatory changes, connective proliferation and sulphur granules, which have also been found in infections caused by *Nocardia brasiliensis*, *Streptomyces madurae* and *Staphylococcus aureus*. These granules are yellowish particles and are formed by clumps of filamentous *Actinomyces* surrounded by neutrophils.³

The usual treatment for actinomycosis consists of high and prolonged doses of antibiotherapy (penicillin, erythromycin, tetracycline, doxycycline, minocycline or clindamycin).⁶ However, in some cases it is necessary to proceed to surgical drainage. In those patients, the duration of antibiotherapy may be reduced (3 months).²

Ethical disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare.

Conflitos de interesse: Os autores declaram não possuir conflitos de interesse.

Financing Support: This work has not received any contribution, grant or scholarship.

Suporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Confidencialidade dos dados: Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Heo SH, Shin SS, Kim JW, Lim HS, Seon HJ, Jung S-I, et al. Imaging of actinomycosis in various organs: a comprehensive review. *RadioGraphics*. 2014;34:19-33.
2. García-García A, Ramírez-Durán N, Sandoval-Trujillo H, Romero-Figueroa MDS. Pelvic actinomycosis. *Canadian Journal of Infectious Diseases and Medical Microbiology*. 2017;2017:1-17.
3. Pérez-López F. Female pelvic actinomycosis and intrauterine contraceptive devices. *Open Access Journal of Contraception* 2010;35.
4. Kim SH, Kim SH, Yang DM, Kim KA. Unusual causes of tubo-ovarian abscess: CT and MR imaging findings. *RadioGraphics*. 2004;24:1575-89.
5. Pusiol T, Morichetti D, Pedrazzani C, Ricci F. Abdominal-pelvic actinomycosis mimicking malignant neoplasm. *Infectious Diseases in Obstetrics and Gynecology*. 2011;2011:1-4.
6. Bennett JE, Dolin R, Blaser MJ. Mandell, Douglas, and Bennett's principles and practice of infectious diseases. (Eight edition). Philadelphia, PA: Elsevier. 2014.