

## Radiological Case Report / Caso Clínico

## Primary Breast Lymphoma – Case Report

*Linfoma Primário da Mama – Caso Clínico*Tatiana de Melo<sup>1</sup>, Elisa Melo Abreu<sup>2</sup><sup>1</sup>Serviço de Imagiologia, Hospital Espírito Santo de Évora, Évora, Portugal<sup>2</sup>Serviço de Radiologia, Centro Hospitalar e Universitário do Porto, Porto, Portugal

## Address

Tatiana de Melo  
Largo do Sr. da Pobreza  
7000-811 Évora, Portugal  
e-mail: tatianafmmelo@gmail.com

Received: 06/10/2021

Accepted: 17/12/2021

Published: 29/04/2022

© Author(s) (or their employer(s)) and ARP  
2022. Re-use permitted under CC BY-NC. No  
commercial re-use.

## Abstract

Breast lymphoma is an uncommon disease and a defying diagnose to establish, both clinically and radiologically. It can be mammographic and sonographically indistinguishable from other breast lesions, both malignant and benign. There are, however, some features that, in conjunction, may suggest this diagnose. We present a clinical case of a primary breast lymphoma as a starting point to review the most common clinical and imagiologic findings of this disease. Since histologically features are relevant to understand and explain the imaging findings, we also exploit the radiologic-pathologic correlation in breast lymphoma. Both staging and treatment of breast lymphoma are very different from other neoplastic breast lesions, so we also emphasize these points. Even though the final diagnose relies on biopsy, it is important to bear in mind the clinical and imaging characteristics that should prompt the diagnose of breast lymphoma, to best evaluate these patients.

## Key-words

Breast; Lymphoma; Mammography;  
Ultrasound; Histology.

## Resumo

O linfoma mamário é raro e o seu diagnóstico desafiante, quer clínica quer imagiologicamente. Nos estudos de mamografia e ecografia mamária, o linfoma mamário pode ser indistinto de outras lesões mamárias, quer benignas quer malignas. Existem, contudo, algumas características que, em conjunto, sugerem este diagnóstico. Apresentamos um caso clínico de linfoma primário da mama como ponto de partida para rever as características clínicas e imagiológicas da doença. As características histológicas são importantes para compreender e explicar os achados imagiológicos, pelo que exploramos a correlação radio-patológica. Tanto o estadiamento como o tratamento do linfoma mamário variam significativamente em comparação com outras lesões neoplásicas mamárias. Apesar de o diagnóstico definitivo ser histológico, consideramos importante saber reconhecer as características clínicas e imagiológicas que devem fazer suspeitar deste diagnóstico, de modo a melhor avaliar estes doentes.

## Palavras-chave

Mama; Linfoma; Mamografia; Ecografia;  
Histologia.

Breast lymphoma (BL) is an uncommon disease, with a prevalence inferior to 0,7%.<sup>1</sup> Furthermore, since there are no specific clinical or imaging findings, BL may be difficult to diagnose both clinically and radiologically.<sup>2</sup> To be able to consider the diagnosis, it is important to be familiar with the clinical and imaging findings that, although not specific for the disease, may arise suspicion. We present a clinical case of a primary breast lymphoma to review this rare pathology, exploiting the radio-pathologic correlations. A 92-year-old female patient, without oncologic or hematologic known disease, presented to the breast unit with a large painful right breast mass, noticed for the first time 6 weeks before. The patient had no other complaints, including no other systemic symptoms. She was chronically medicated with antihypertensive, oral antidiabetics and oral anticoagulant drugs.

Clinical evaluation revealed a firm oval mass at the transition of outer quadrants (TOQ), measuring about 7cm. No other masses or nipple discharge were present, bilaterally. No axillary lymphadenopathies were palpable.

Mammographic evaluation revealed a global asymmetry of the right breast caused by a high-density oval lesion at the TOQ with obscured margins. There was concomitant

diffuse skin thickening (Figure 1). No clusters of suspicious calcifications were present.


Ultrasound revealed a single oval mass, with indistinct margins and very heterogenous echo pattern, with echogenic and anechogenic areas. The long axis of the mass was parallel to the skin line and it was highly vascularized (Figure 2). No axillary lymphadenopathies were seen.

Giving these findings and considering the absence of other complaints, we suspected of a primary breast malignancy. Our first diagnostic options were a poorly differentiated carcinoma, invasive lobular carcinoma or even a breast lymphoma.

To achieve diagnosis, we performed a 14G core needle biopsy of the lesion. The pathologic analysis revealed a large B-cell lymphoma.

Thoraco-abdominal-pelvic computed tomography (CT) was performed to evaluate systemic disease, revealing a large homogenous and well circumscribed lesion at the outer quadrants of the right breast (Figure 3). There weren't any lymphadenopathies, neither organomegalies.

Patient was proposed to chemotherapy, with good response. BL can be either primary or secondary, and like other lymphomas, has a bimodal age distribution with peaks in the middle of the 4th and 7th decades of life.<sup>2,3,4</sup>




**Figure 1** – Mammographic evaluation with standard projections [(A) mediolateral oblique view right breast; (B) mediolateral oblique view left breast; (C) craniocaudal view right breast; (D) craniocaudal view left breast] shows a global asymmetry caused by a high-density, non-calcified, oval lesion at the TOQ of the right breast, with indistinct and irregular margins. Diffuse skin thickening of the right breast is present.

According to Wiseman and Liao, the diagnose of primary breast lymphoma (PBL) requires an adequate biopsy demonstrating close anatomic proximity of breast parenchyma and lymphomatous infiltrate, absent history of previous extramammary lymphoma, and no evidence of concurrent widespread disease, except for simultaneously appearing ipsilateral axillary lymph nodes.<sup>5</sup> Most PBL are intermediate or high-grade lymphomas, frequently B-cell lineage and, rarely, T-cell or histiocytic lineages.<sup>3,6</sup> Hodgkin's disease occurs very rarely as a primary tumour of the breast.<sup>7</sup>


Thus, secondary breast lymphoma (SBL) is diagnosed in the presence of widespread disease as a metastatic lesion, occurring frequently in patients with diffuse large B-cell lymphoma.<sup>1</sup>

There are no significant imaging differences between PBL and SBL, nor pathognomonic imaging findings. So, BL can be mammographic and sonographically indistinguishable from primary breast carcinoma (BC), fibroadenoma, phyllodes tumour or even metastatic disease.<sup>7</sup> Diagnosis relies on adequate tissue biopsy for histopathological evaluation and immunophenotyping.<sup>8</sup>



**Figure 2** – Breast ultrasound reveals (A) a large single oval mass, with indistinct margins and a heterogenous echo pattern, with echogenic and anecogenic areas. (B) Because of its indistinct margins and large dimension, it was difficult to accurately measure the lesion at sonography, and it was underestimated. (C) The evaluation with colour Doppler shows prominent arteries and veins at the periphery of the lesion.

PBL frequently presents as a large or rapidly enlarging solitary painless mass, especially in younger patients, clinically mimicking BC.<sup>2</sup> On the other hand, multifocal, multicentric or bilateral lesions suggest SBL. Alternatively, BL may present as a painful mass or as breast skin thickening, especially those of high grade.<sup>2,3,5</sup> Most cases of PBL do not have prominent axillary lymphadenopathy, although this can be present.<sup>2</sup> Systemic B-symptoms are infrequent in PBL, but common in SBL.<sup>1,3</sup> Histologically features are relevant to understand the imaging findings. The pathological examination shows



**Figure 3** – Computed tomography images [(A) axial; (B) coronal] demonstrate the mammary lesion as a large homogenous lesion at the outer quadrants of the right breast (circle).

lymphoma cells infiltrating the breast lobules and ducts, which remain intact and do not show neoplastic changes.<sup>5</sup> BL are highly cellular lesions, composed of non-cohesive round lymphoid cells of variable sizes, closely packed, with small extracellular spaces.<sup>9</sup> No desmoplastic or fibrotic reaction is usually present.<sup>9,10</sup> Lymphangitic spread can occur, causing overlying skin changes.<sup>11</sup>

On mammograms, BL most commonly occurs as a solitary oval or round mass of homogeneous density, uncalcified (Figure 1). Since fibrotic or desmoplastic reaction are absent, BL does not cause stromal distortion of the surrounding breast parenchyma, contrary to BC.<sup>2,3,7,10</sup> BL can also be mammographically occult or present as a diffuse increased parenchyma density.<sup>7,10</sup> Skin thickening can be appreciated on mammograms (Figure 1). This last pattern is predominantly observed with high-grade lymphoma.<sup>10</sup> The findings of spiculate borders or calcification should propound another diagnose.<sup>7</sup>

Ultrasound features are nonspecific, so a wide spectrum of appearances have been described.<sup>2,3</sup> Some authors have reported a common presentation as a solitary hypervascular mass (Figure 2), with irregular shape and indistinct margins, homogenous hypoechoic or a heterogeneous (Figure 2) echo pattern, without posterior acoustic phenomena.<sup>2,3</sup>

MRI features are also nonspecific, often like those of BC.<sup>1,5</sup> A round or oval mass, with areas of hypointensity or isointensity at T1-weighted imaging (WI), and areas of hyperintensity at T2-WI. The enhancement pattern may be homogeneous or heterogeneous, most commonly with a type II kinetic curve or, less frequently, a type III kinetic curve.<sup>1,5</sup> Skin thickening is more apparent than at mammography. The essential role of MRI is to evaluate the presence and extent of multicentric and multifocal lesions and stage the disease.<sup>1</sup> It is also useful to evaluate treatment efficacy, and diagnose recurrent disease.<sup>1</sup> Since these are highly cellular lesions, with small extracellular spaces, they often demonstrate higher restriction and lower ADC values than BC.<sup>9</sup>

CT is an important tool to stage, restage, and evaluate treatment response, being the modality of choice if Positron Emission Tomography (PET)-CT is not available (Figure 3). As others glucose-avid breast cancers, BL demonstrates high uptake at fluorodeoxyglucose (FDG) at PET-CT. Similarly, to other lymphomas, the treatment of BL is based on chemotherapy and radiation.<sup>12</sup> Surgical treatment with lumpectomy is an option in some cases.

#### Ethical disclosures / Divulgações Éticas

*Conflicts of interest:* The authors have no conflicts of interest to declare.

*Conflitos de interesse:* Os autores declaram não possuir conflitos de interesse.

*Financing Support:* This work has not received any contribution, grant or scholarship.

*Suporte financeiro:* O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

*Confidentiality of data:* The authors declare that they have followed the protocols of their work center on the publication of data from patients.

*Confidencialidade dos dados:* Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

*Protection of human and animal subjects:* The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

*Proteção de pessoas e animais:* Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

#### References

- Raj SD, Shurafa M, Shah Z, Raj KM, Fishman MDC, Dialani VM. Primary and secondary breast lymphoma: clinical, pathologic, and multimodality imaging review. *Radiographics*. 2019;39:610-25.
- Lyou CY, Yang SK, Choe DH, Lee BH, Kim KH. Mammographic and sonographic findings of primary breast lymphoma. *Clin Imaging*. 2007;31:234-8.

- Yang WT, Lane DL, Le-Petross HT, Abruzzo L V, Macapinlac HA. Breast lymphoma: imaging findings of 32 tumors in 27 patients. *Radiology*. 2007;245:692-702.
- Jeon HJ, Akagi T, Hoshida Y, et al. Primary non-hodgkin malignant lymphoma of the breast: an immunohistochemical study of seven patients and literature review of 152 patients with breast lymphoma in Japan. *Cancer*. 1992;70:2451-9.
- Wiseman C, Liao KT. Primary lymphoma of the breast. *Cancer*. 1972;29:1705-12.
- Arber DA, Simpson JF, Weiss LM, Rappaport H. Non-Hodgkin's Lymphoma Involving the Breast. *Am J Surg Pathol*. 1994;18:288-95.
- Lieberman L, Deutch BM, Giess S, Louie C. Non-Hodgkin lymphoma of the breast and correlation with histopathologic findings. *Radiology*. 1994;192:157-60.
- Mason HS, Johari V, March DE, Crisi GM. Primary breast lymphoma: radiologic and pathologic findings. *Breast J*. 2005;11:495-6.
- Zhou C, Lv K, Lin D, Yu Y, Xu M, Bu Y. Radiological analysis of breast lymphoma: experiences from cases series studies. *Med (United States)*. 2019;98:e18101.
- Sabaté JM, Gómez A, Torrubia S, et al. Lymphoma of the breast: clinical and radiologic features with pathologic correlation in 28 patients. *Breast J*. 2002;8:294-304.
- Kalli S, Lanfranchi M, Alexander A, Makim S, Freer PE. Spectrum of extramammary malignant neoplasms in the breast with radiologic-pathologic correlation. *Curr Probl Diagn Radiol*. 2016;45:392-401.
- Nicholson BT, Bhatti RM, Glassman L. Extranodal lymphoma of the breast. *Radiol Clin North Am*. 2016;54:711-26.