

Radiological Case Report / Caso Clínico

Cervical Haematoma Secondary to Spontaneous Thyroid Haemorrhage

Hematoma Cervical Secundário a Hemorragia Espontânea da Tireoide

Matilde Almeida e Sousa¹, Eduardo Dutra², Guida Matos Ferreira¹

¹Radiology Department, Centro Hospitalar Universitário Lisboa Central, Lisboa, Portugal
²Pathology Department, Centro Hospitalar Universitário Lisboa Central, Lisboa, Portugal

Address

Matilde Almeida e Sousa
Radiology Department
Centro Hospitalar Universitário Lisboa Central
Rua da Beneficência, 8
1050-099 Lisboa, Portugal
e-mail: sousamatilde@campus.ul.pt

Received: 16/11/2021**Accepted:** 20/12/2021**Published:** 29/04/2022

© Author(s) (or their employer(s)) and ARP
2022. Re-use permitted under CC BY-NC. No commercial re-use.

Abstract

Spontaneous thyroid haemorrhage is rare, but it can be a potentially life-threatening condition in severe cases, due to acute upper airway compromise.

We report the case of a 75-year-old patient on chronic anticoagulation therapy for atrial fibrillation, who presented with sudden painful neck bulging and dysphagia. Physical examination revealed a hard mass on the anterior left side of the neck. Preliminary laryngoscopy was normal. The diagnosis was made using ultrasonography and further characterization with computed tomography, showing haemorrhage of a thyroid nodule with capsular disruption and diffusion of the haematoma to the deep cervical spaces, causing important tracheal deviation and hypopharynx compression. The patient developed dyspnoea and ultimately required urgent hemithyroidectomy.

This case also highlights the importance of having a high index of clinical suspicion and prompt imaging in patients with atypical presentations of a neck mass.

Key-words

Thyroid gland; Thyroid nodule;
Thyroidectomy; Haematoma; Neck.

Resumo

A hemorragia espontânea da tireoide é rara, mas pode ser uma condição potencialmente fatal nos casos graves, por compromisso agudo da permeabilidade da via aérea superior.

Apresentamos o caso clínico de uma mulher de 75 anos cronicamente anticoagulada, por fibrilhação auricular, que recorre ao Serviço de Urgência por tumefação cervical dolorosa de aparecimento súbito, com disfagia. Ao exame objetivo apresentava uma massa dura e dolorosa na vertente anterior paramediana esquerda do pescoço. A laringoscopia preliminar foi normal. O diagnóstico foi feito por ecografia e posterior caracterização com tomografia computadorizada, que revelaram hemorragia de nódulo tireoideu, com disrupção capsular e extensão do hematoma para os espaços cervicais profundos, causando importante desvio traqueal e compressão da hipofaringe. Pelo agravamento clínico com desenvolvimento de dispneia, procedeu-se a hemitiroidectomia urgente.

Este caso destaca a importância de um elevado índice de suspeição clínica e avaliação imagiológica atempada em doentes com apresentações atípicas de massa cervical.

Palavras-chave

Glândula tireoide; Nódulo da tireoide;
Tiroidectomia; Hematoma; Pescoço.

Introduction

Although highly vascularised and with a significant incidence of benign and malignant diseases that further increase its vascularisation, spontaneous thyroid bleeding is rare, and most cases are mild.^{1,2,3} However, the risk of bleeding due to goitre or nodular disease-induced abnormal vascularity, triggered by the use of anticoagulants, can cause massive haemorrhage and sudden unexpected airway compromise.^{1,2,4}

We describe a case of spontaneous bleeding of a thyroid nodule, with capsular disruption and extension of the haematoma to the deep cervical planes, in a chronic anticoagulated patient.

Case Presentation

A 75-year-old woman presented to the emergency department complaining of a painful mass on the left side of the neck that developed overnight, along with dysphagia. There was no history of trauma, fever, cough, recent medical procedures or previous thyroid disease.

She had a known background of hypertension, polycythaemia vera and atrial fibrillation and was medicated with hydroxyurea, lercanidipine, amiodarone, clopidogrel, olmesartan and tinzaparin sodium, a low molecular weight heparin.

The patient was afebrile, eupnoeic at rest, without stridor. Neck examination revealed a hard lesion confined to the left side of the neck with approximately 10 cm, painful on palpation, adherent to surrounding tissues, non-pulsatile, without associated palpable adenopathies in the cervical chains. Preliminary laryngoscopy was otherwise normal, except for paralysis of the left vocal cord.

Laboratory workup showed a slightly low haemoglobin level of 11.1g/dL [normal range = 12.1 to 15.1 g/dL], platelet count of 522×10^9 [normal range = $150-450 \times 10^9$], normal prothrombin time of 12.9s, elevated activated partial thromboplastin time of 58.1 s [normal range = 25.1-36.5s] and elevated fT4 of 1.54 g/dL [normal range = 0.54-1.24 g/dL] with normal thyroid-stimulating hormone (TSH) level.

Ultrasound study of the neck (Fig. 1) showed a left lobe thyroid nodule measuring approximately 85x60x55mm,



Figure 1 – Cervical ultrasound (A-C) demonstrates a thyroid nodule with mixed cystic and solid content, central inhomogeneous fluid, fluid-fluid level and trachea (T) deviation.

with mixed cystic and solid content, with central inhomogeneous fluid and fluid-fluid level, causing trachea deviation, suggestive of a haemorrhagic thyroid nodule. To establish the extension of the process, a neck Computed Tomography (CT) was obtained (Fig. 2), without contrast administration because of significant renal impairment, confirming a large thyroid nodule in the left lobe, with areas of recent haemorrhage depicted by intra-nodular hyper-attenuating portions and fluid-fluid levels, with apparent posterosuperior disruption of the capsule and extension of the process to the hypopharynx with mass effect, causing obliteration of the pyriform sinus and deviation of the trachea. There was extensive oedema in the left parapharyngeal space, carotid space and crossing the midline through the retropharyngeal/danger spaces. Indirect laryngoscopy was repeated and revealed a submucosal left hypopharynx haematoma with obliteration of the pyriform sinus and permeable glottis.

to the pharynx/hypopharynx and laterosuperior cervical planes, with significant tracheal deviation. The study of the surgical specimen (Fig. 3) confirmed the existence of a dominant hyperplastic nodule in the left lobe, with extensive central haemorrhage and pseudocapsule rupture. The bleeding extended to the adjacent soft tissues. There were no neoplastic cells. The patient clinically improved and was discharged five days after surgery.

Discussion

Although the thyroid is a very vascularized organ, spontaneous haemorrhage, not related to previous trauma or interventional procedures, is not frequent and usually confined to the gland. There are not many cases described in the literature.




Figure 2 – Non-contrast-enhanced soft-tissue window neck CT scan, with multiplanar reformats on the axial (A,C) and coronal (B) planes. A and B show a large thyroid nodule in the left lobe with intra-nodular hyper-attenuating portions and a fluid-fluid level, with apparent disruption of the capsule posteriorly (arrow), and significant tracheal deviation. The axial plane of the hypopharynx (C) shows the extension of the bleeding outside the thyroid, with obliteration of the pyriform sinus and the left parapharyngeal, carotid and retropharyngeal/danger spaces.




Figure 3 – A and B Haematoxylin and Eosin staining - Low power view (x40) showing a dominant thyroid nodule composed of variable size follicles, with extensive intra-nodular haemorrhage (star) extending to the surrounding tissues (arrow).

Based on the clinical and radiological findings, spontaneous thyroid nodule bleeding was diagnosed with capsular disruption and hypopharynx haematoma. Anticoagulation was promptly reversed. The patient developed respiratory distress, therefore, after securing the airway, she underwent urgent hemithyroidectomy. The surgical exploration described a left lobe thyroid nodule with extensive haemorrhage and capsular rupture on its superior part, with an extension of the haematoma

Pre-existing thyroid conditions such as solitary or multiple benign or malignant nodules, cystic nodules and colloid goitre are risk factors increasing the vascularity of the gland and vulnerability for bleeding.^{1,2,3,4} The origin of the haemorrhage is mainly venous, and the possible mechanisms may be related to abnormal thyroid vascular anatomy with venous fragility and the creation of arteriovenous shunting inside the nodules, conditioning intra-nodular haemorrhage. Factors favouring

an increase in venous pressure, such as Valsalva, coughing or exercise, may raise the probability of spontaneous bleeding.^{2,4,5,6} Moreover, patients on anticoagulation therapy have more risk of haemorrhage, although no cases of thyroid haemorrhage were revealed in the trials across all anticoagulants, maybe because this complication is not captured in many of these trials.^{1,2}

In mild cases, patients with acute thyroid bleeding present sudden neck bulging with local discomfort or pain and frequent dysphagia. In severe cases, it can result in a large expanding haematoma with capsule rupture, with haemorrhage extending to cervical planes, resulting in sudden compression of the trachea, causing dyspnoea and stridor, and leading to possible upper airway obstruction.^{1,2,4,5} Furthermore, thyroid bleeding can cause a sudden release of preformed thyroid hormones and acute hyperthyroidism, which is usually self-limiting, needing no treatment.¹ It is essential to consider other possible and rare causes of spontaneous cervical haemorrhage, such as lesions of the upper and lower thyroid arteries and lesions of the parathyroid glands.⁷

Laboratory blood tests for coagulation and hormonal status should be obtained. Ultrasound evaluation is essential for proper diagnosis and may be helpful in distinction from other acute neck conditions. Nevertheless, in more severe cases, when there is massive bleeding with clinical signs of airway or hemodynamic compromise, patients

should be evaluated by CT scan, preferably with contrast administration.⁴

As a therapeutic approach, securing the airway with endotracheal intubation is the most critical action before any diagnostic and treatment procedures, and, if possible, anticoagulation should be reversed immediately.² Conservative treatment is the preferred choice for most patients, but early surgical decompression should be considered whenever tracheal compression is a concern or when bleeding is not controlled.^{1,5} Patients managed conservatively should be referred for ambulatory endocrinology follow-up, namely because bleeding could arise as the first sign of malignant thyroid disease.⁴

Conclusion

It is essential to remember that thyroid haemorrhage can have life-threatening presentations, severe enough to require emergency surgery. Knowing the typical clinical signs and symptoms is mandatory in order to have a good and fast approach. Imaging features of thyroid bleeding can ensure the proper diagnosis and lead to adequate and prompt treatment. Therefore, the collaboration between emergency department physicians, otorhinolaryngologists, radiologists and general surgeons is needed for a satisfactory outcome in these cases.

Ethical disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare.

Conflitos de interesse: Os autores declaram não possuir conflitos de interesse.

Financing Support: This work has not received any contribution, grant or scholarship.

Suporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Confidencialidade dos dados: Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Kokatnur L, Rudrappa M, Mittadodla P. Acute airway obstruction due to spontaneous intrathyroid hemorrhage precipitated by anticoagulation therapy. *Indian J Crit Care Med.* 2014;18:825-7.
2. Gunasekaran K, Rudd KM, Murthi S, Kaatz S, Lone N. Spontaneous thyroid hemorrhage on chronic anticoagulation therapy. *Clinics and Practice.* 2017;7:932.
3. González-Cruz A, García-Ferrer L, García C. Cervical and mediastinum haematoma secondary to spontaneous thyroid rupture. *Acta Otorrinolaringol Esp.* 2010;61:459-61.
4. Covino M, Princi P, De Luca G, Ciello A, Simeoni B, Boanchi A, et al. Spontaneous thyroid nodule hemorrhage in the emergency department. *Endocr Pract.* 2020;26:192-6.
5. Lee JK, Lee DH, Cho SW, Lim SC. Acute airway obstruction by spontaneous hemorrhage into thyroid nodule. *Indian J Otolaryngol Head Neck Surg.* 2011;63:387-9.
6. Vijapurapu R, Kaur K, Crooks, NH. A case of airway obstruction secondary to acute haemorrhage into a benign thyroid cyst. *Case Reports in Critical Care.* 2014; Article ID 372369.
7. Petersen NK, Udholm S, Londero SC. Acute airway obstruction due to spontaneous intrathyroidal haemorrhage. *BMJ Case Rep.* 2021;14:e238935