

Images of Interest / Imagens de Interesse

Vaginal Lithiasis – A Rare Case*Litíase Vaginal – Um Caso Raro*Cristina Mota¹, Inês Dias Marques¹, Ana Teresa Almeida¹¹Serviço de Imagiologia, Centro Hospitalar de Vila Nova de Gaia / Espinho, EPE, Portugal**Address**

Cristina Mota
Serviço de Imagiologia
Centro Hospitalar de Vila Nova de Gaia /
Espinho, EPE
R. Conceição Fernandes
4434-502 Vila Nova de Gaia, Portugal
e-mail: cristina.sousa.mota@chvng.min-saude.pt

Received: 03/04/2022

Accepted: 21/12/2022

Published: 27/09/2023

© Author(s) (or their employer(s)) and ARP
2023. Re-use permitted under CC BY-NC. No
commercial re-use.

Abstract

Vaginal lithiasis is a rare pathology often misdiagnosed due to low diagnostic suspicion. We present a case of an 83-year-old female with primary vaginal lithiasis.

Keywords

Vaginal lithiasis; Urinary stasis; Dysuria.

Resumo

Litíase vaginal é uma patologia rara, diagnosticada poucas vezes devido à baixa suspeição diagnóstica. Apresentamos o caso de uma mulher de 83 anos com litíase vaginal primária.

Palavras-chave

Litíase vaginal; Estase urinária; Disúria.

Case

A dependent 83-year-old female presented to the emergency department with a history of dysuria and prostration. The patient had a past medical history of cervical cancer, treated with chemotherapy and radiotherapy, which precipitated enterovesical and enterovaginal fistulas. No abnormal findings were noticed on physical examination. An abdominal and pelvic computed tomography (CT) with intravenous contrast was performed which revealed the presence of multiple calcifications within the vagina (Figure 1), surrounded by a big amount of gas, without clear evidence of vesicovaginal or urethrovaginal fistula. These findings were suggestive of primary vaginal lithiasis. Stones were also observed within the bladder (Figure 1).

Discussion

Vaginal lithiasis, also known as vaginal stones or calculi, is a rare pathology that is often misdiagnosed because most clinicians do not even suspect its presence.

The formation of vaginal stones is a slow process, and most cases are only discovered when the stones are large enough to cause obvious clinical symptoms, or accidentally when an imaging exam is performed.¹

Clinically, the patients could present dysuria, vaginal pain, and dyspareunia. Additionally, vaginal calculi can also cause urinary tract irritation symptoms such as frequent urination and urination urgency.¹

Based on the etiology of the disease, there are 2 types of vaginal stones: primary and secondary. Primary vaginal



Figure 1 a,b – Vaginal lithiasis. Sagittal A) and axial B) images from contrast CT show multiple calculi in the vagina and the bladder.

lithiasis results from the stasis of urine within the vagina. Stagnant urine facilitates infections caused by urease-producing bacteria, such as *Klebsiella*, *Proteus mirabilis*, or *Escherichia coli*, so that the acidic environment of the vagina becomes alkaline, which contributes to the formation of vaginal stones. Primary vaginal lithiasis is seen in cases of congenital anomalies of the genitourinary tract, vesicovaginal or urethrovaginal fistula, trauma, ectopic ureters, previous pelvic radiotherapy, neuropathic bladder, and other causes of vaginal outlet obstruction.^{1,2} Secondary vaginal stones result from the crystallization of urinary constituents around foreign bodies introduced into the vagina, including retained medical gauze, missed vaginal pessary, or an intrauterine contraceptive device.²

In patients suspected of having vaginal lithiasis, ultrasound and X-ray examinations of the pelvis are helpful in the establishment of the correct diagnosis. Ultrasonography can confirm the presence of stones and allow their localization in the pelvic organs.^{1,2} When the stones are difficult to identify,

it is feasible to perform a pelvic CT and magnetic resonance imaging for the correct diagnosis. In our particular case, the diagnosis was performed with the aid of a pelvic CT.

Treatment involves the complete removal of vaginal stones, either through manual extraction or surgery. Surgery is the preferred treatment, however, stone fragmentation and extracorporeal shock wave lithotripsy, as well as endoscopic intervention, are commonly employed techniques.³ Recurrence is only avoided if the underlying etiology is adequately addressed.

Conclusion

The diagnosis of vaginal lithiasis can be difficult and requires that the physician is highly suspicious of this possible diagnosis. As in our case, this diagnosis should be considered when urinary symptoms are present in women with physical disabilities, particularly those who are bedridden and have urinary incontinence.

Ethical Disclosures / Divulgações Éticas

Conflicts of interest: The authors have no conflicts of interest to declare.

Conflitos de interesse: Os autores declaram não possuir conflitos de interesse.

Financing Support: This work has not received any contribution, grant or scholarship.

Suporte financeiro: O presente trabalho não foi suportado por nenhum subsídio ou bolsa.

Confidentiality of data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Confidencialidade dos dados: Os autores declaram ter seguido os protocolos do seu centro de trabalho acerca da publicação dos dados de doentes.

Protection of human and animal subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki).

Proteção de pessoas e animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pelos responsáveis da Comissão de Investigação Clínica e Ética e de acordo com a Declaração de Helsínquia da Associação Médica Mundial.

References

1. Wei D, Xie Y, Niu X. Vaginal stones caused by urethrovaginal fistula: a case report. *Medicine*. 2019;98:e18003.
2. Castellan P, Nicolai M, De Francesco P, Di Tizio L, Castellucci R, Bada M, et al. Primary vaginal calculus in a woman with disability: case report and literature review. *J Endourol Case Rep*. 2017;3:182-5.
3. Jaspers JW, Kuppens SM, van Zundert AA, de Wildt MJ. Vaginal stones in a 5-year-old girl: a novel approach of removal. *J Pediatr Adolesc Gynecol*. 2010;23:e23-5.