

CASO CLÍNICO

Cervical Blood Patch: A Case Report of an Unusual Technique for an Unusual Diagnosis

Blood Patch Cervical: Uma Técnica Inabitual para um Diagnóstico Inabitual

José Cravo^{1*} , Ionut Tabolcea¹ , Turgay Tuna¹ 

Afiliação

¹ Cliniques Universitaires de Bruxelles – Hôpital Erasme, Bruxelas, Bélgica.

Keywords

Anesthesiology ; Blood Patch, Epidural; Cerebrospinal Fluid Leak

Palavras-chave

Placa de Sangue Epidural ; Vazamento de Líquido Cefalorraquidiano

ABSTRACT

Cervical epidural blood patches (CEBP) remain an unfamiliar domain for most anesthesiologists as they fear the possible neurological complications related with the procedure. The injection of autologous blood as close as possible to the dural leakage site has shown to be the most effective approach for symptom resolution in case of spontaneous cervical dural tears. This case report describes in detail an alternative and feasible technique to perform an epidural blood patch at the cervical level.

RESUMO

A realização de *blood patches* cervicais continuam a ser um terreno inusual para a maioria dos anestesistas. Este facto deve-se principalmente à falta de experiência nesta técnica e ao receio de possíveis complicações neurológicas associadas a este procedimento. A injeção de sangue autólogo o mais perto possível do ponto de fuga de líquido cefalorraquidiano demonstra ser a técnica mais eficaz para a resolução de sintomas resultantes de uma fuga espontânea a nível cervical. Este caso clínico descreve em detalhe uma técnica alternativa e factível para a realização de um *blood patch* cervical, de forma a minimizar o risco teórico associado a este procedimento.

INTRODUCTION

Intracranial hypotension headache is almost always related to a cerebrospinal fluid (CSF) leakage due to a dural tear, which can have an iatrogenic, traumatic or spontaneous etiology. The incidence of spontaneous dural tears is around 5 in 100 000 individuals per year¹ and it is more common in women (2:1 ratio). Its etiology and pathophysiology are unknown and it remains a diagnosis of exclusion, when there is no previous history of trauma, spinal surgery, lumbar puncture nor an epidural neuroaxial procedure. Patients with spontaneous intracranial hypotension typically present with postural headache, increasing in intensity in a sitting or upright position and achieving a significant headache relieve by laying down in a horizontal position, explained by the effect of gravity in the amount of CSF surrounding the brain. In some cases, other neurological symptoms can be present, such as nausea, vomiting, photophobia, blurred vision, diplopia, tinnitus, dysgeusia, among others. Diagnosis is based on clinical history and symptoms, and confirmed by brain and spine magnetic resonance imaging (MRI) where indirect findings of a dural leak can be seen, such as the curtain sign, when an extradural fluid collection separates the dura from the bony spinal canal,² but also the enhancement of pachymeninges, pituitary hyperemia, engorgement of venous structures and/or sagging of the brain. Myelography can be helpful in identifying the precise site of CSF leakage. Initial management includes conservative measures based on oral analgesics and fluid hydration. In refractory patients, an invasive treatment technique may be required, known as 'blood patch', characterized by the injection of autologous blood in the epidural space.

In this patient's case, we report a feasible and effective cervical blood patch technique, based on the utilization of an epidural catheter introduced at a lower cervical level, to reach a cervical CSF leakage at a higher cervical level, where

Autor Correspondente/Corresponding Author*:

José Francisco Cravo

Morada: Route de Lennik 808, 1070 Bruxelas, Bélgica.

E-mail: francisco.cravo@erasme.ulb.ac.be

the risk of complications of a cervical epidural blood patch (CEBP) would be higher.

This case report was approved by the local Ethics Committee with the reference code P2020/644, patient's written informed consent was obtained and signed, and EQUATOR's-enhancing the quality and transparency off health research guidelines were followed in its elaboration.

CASE REPORT

A 35-year-old female patient presents to a Pain Management Clinic consultation on a secondary center, referred to by her General Practitioner, with a proposal of bilateral infiltration of the greater and lesser occipital nerves in relation with a 4 weeks lasting bilateral headache. The patient describes her pain as an intense tight band around the head, affecting as well her cervical and occipital regions. The pain appeared at the first rise in the morning and significantly decreased in supine position. The palpation of the occipital and cervical region did not alter the pain's character nor intensity. The patient had no relevant past medical history nor history of recent trauma. This clinical description led the attending physician to cancel the planned infiltration and proceed to further diagnostic evaluation with the realization of a head and total spine magnetic resonance imaging (MRI), leading to the diagnosis of a cervical dural tear with an adjacent extradural liquid collection at C2C3 level. The MRI showed also signs of intracranial hypotension like hypophyseal hyperemia, venous plexus engorgement and pachymeningeal enhancement. Conservative treatment was instaured with oral hydration and analgesics, and the patient was referred to a tertiary center capacitated to perform cervical blood patches.

The CEBP was performed by a team composed of two experienced anesthesiologists, an anesthesiology trainee and a nurse. The procedure's details and its potential risks were explained to the patient and she was asked to fast for 6 hours before the procedure. An intravenous line was placed upon arrival in the operating room. The patient was placed in prone position on the operating table and the fluoroscope positioned in order to allow the team to easily obtain both an anteroposterior and a profile views. The posterior aspect of the cervical region was disinfected with a 2% chlorhexidine solution. Under local anesthesia with lidocaine 2%, a 17G Tuohy needle was introduced into the epidural space at the C5C6 level (Fig. 1), under fluoroscopy guidance and using the hanging drop technique. An epidural catheter was introduced still under fluoroscopic control, with the catheter tip reaching the C2C3 level, where the dural tear was detected in the diagnostic imaging studies. There was no blood nor cerebrospinal fluid upon aspiration testing. The catheter tip position was confirmed by the injection of a total of 2 mL of contrast solution (Fig. 2). The Tuohy needle was




Figure 1. Fluoroscopy profile view of the introduction of a 17G Tuohy needle into the epidural space, at the C5C6 level




Figure 2. Contrast product was injected through an epidural catheter confirming its position at the C2C3 level




Figure 3. After removal of the Tuohy needle, the epidural catheter remained in place, at the C2C3 level. Diffused contrast product remains visible in the epidural space

then removed, leaving the epidural catheter in place (Fig. 3) while the team proceeded to a sterile blood withdraw from a peripheral vein, and a total of 10 mL were injected through the epidural cervical catheter. The patient remained under surveillance on the recovering room for 4 hours, in a strict dorsal decubitus. There were no complications related with the procedure and neurological examination before discharge was strictly normal. The patient's follow-up consisted on a telephonic consultation one week after the procedure and a standard pain clinic consultation 20 days later, where the patient complained of remaining mild cervicalgia but the typical symptoms of intracranial hypotension headache were no longer present. A follow-up cervical spine MRI was performed and it resulted completely normal, showing no signs of intracerebral hypotension, cervical dural tear nor extradural or paravertebral fluid collections.

DISCUSSION

Spontaneous CSF leakages occur most commonly at the cervicothoracic spine, but lumbar leakages have been described.³ Although some reports showed that a lumbar EBP may lead to a partial, and sometimes, even complete symptom resolution independently of the leakage origin,⁴ literature demonstrates that an EBP close to the site of CSF leak is more effective.⁵ Currently accepted hypothesis suggest that the increase in hydrostatic pressure resulting from the acute compression of the thecal sac might provide an immediate, partial or complete relief of symptoms, but this effect being brief in duration; while the dural leak sealing by the tamponade effect provided by the blood clot would be responsible for the permanent resolution of symptoms, allowing the gradual restoration of CSF volume. Despite the fact that an epidural blood patch close to the leakage site seems to be more effective, most anesthesiologists are more experienced in lumbar EBP and in lumbar epidural space approach in general, while the cervical approach remains an unfamiliar domain, mainly due to the higher risks involved in relation with the narrower diameter of the cervical epidural space. Possible complications associated with a cervical EBP range from minor complications, such as local pain or radiculopathy, to more severe cases including direct spinal cord injury, spinal cord compression and spinal cord infarction, all of which may lead to paraplegia, tetraplegia and even death.⁶ All the previously mentioned complications, with the exception of local pain at the site of injection, are deemed rare, and to our knowledge, besides a few punctual case reports, there is no reliable statistical data on their incidence and frequency.

Until now, there are no published studies evaluating the exact volume that should be injected when performing a cervical EBP but a literature review by Kapoor *et al*, showed that volumes up to 15 mL of blood have been injected in the

epidural space at the cervical level, and up to 20 mL at the C7-Th1 level, without any serious complications.⁶ Therefore, in this patient's case, a safest technique was chosen, by approaching the cervical epidural space at a lower level (C5C6), where the epidural space diameter is larger than at the upper cervical levels, and an epidural catheter was then introduced under fluoroscopy guidance up to the dural leakage level. Position confirmation was obtained by contrast injection, allowing the delivery of a total volume of 10 mL, the closest to the leakage site as possible, maximizing the efficiency of the procedure.

There are no standard of care guidelines establishing how cervical EBPs should be performed. Regarding the patient's position, the existing reported cases preferred the use of prone position, and in fewer cases the lateral decubitus. In terms of imaging guiding techniques, fluoroscopy was the chosen option in most of the cases described in literature, more than computerized tomography; this results mainly from easier access and availability, as well as higher level of practical experience with the technique. In the reported case, we chose the prone position with slight cervical flexion to promote the theoretical increase in the cervical epidural space of 1 to 2 mm,⁷ and we used fluoroscopic guidance mainly due to prompt availability and higher degree of experience of the operators with this type of imaging technique. Concerning the cervical EBP technique itself, we considered that the patient's safety should remain the top priority, so we preferred a lower level puncture approach plus epidural catheter introduction up to the leakage site, rather than a direct leakage level puncture at an upper cervical level, which would pose a higher risk of procedural complications due to narrower epidural space diameter at this level.

CONCLUSION

Cervical EBP is considered the most effective treatment of refractory hypotension headache secondary to a cervical dural CSF leakage. The number of studies and reported cases in literature related with cervical EBP remain sparse and limited, while procedure related safety and potential complications are the main concerns among anesthesiologists.

This patient's case intends to report the procedural aspects and details related with the performance of a cervical EBP to treat a spontaneous cervical dural tear at C2C3 level, where the use of an epidural catheter introduced at C5C6 level reaching up to the level of the dural tear, allowed to avoid a more risky approach at a higher cervical level.

This case report demonstrates a feasible and effective EBP technique to approach high cervical level dural tears.

Ethical Disclosures

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financing Support: This work has not received any contribution, grant or scholarship.

Confidentiality of Data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Patient Consent: Consent for publication was obtained.

Provenance and Peer Review: Not commissioned; externally peer reviewed.

Responsabilidades Éticas

Conflitos de Interesse: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

Fontes de Financiamento: Não existiram fontes externas de financiamento para a realização deste artigo.

Confidencialidade dos Dados: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

Consentimento: Consentimento do doente para publicação obtido.

Proveniência e Revisão por Pares: Não comissionado; revisão externa por pares.

ORCID

José Francisco  <https://orcid.org/0000-0002-2519-7079>

Ionut Tabolcea  <https://orcid.org/0000-0001-9475-7001>

Turgay Tuna  <https://orcid.org/0000-0003-1032-794X>

Received: 20th of February, 2021 | Submissão: 20 de fevereiro, 2021

Accepted: 13th of May, 2021 | Aceitação: 13 de maio, 2021

Published: 30th of September, 2021 | Publicação: 30 de setembro, 2021

© Author(s) (or their employer(s)) and SPA Journal 2021. Re-use permitted under CC BY-NC. No commercial re-use.

© Autor (es) (ou seu (s) empregador (es)) Revista SPA 2021. Reutilização permitida de acordo com CC BY-NC. Nenhuma reutilização comercial.

REFERENCES

1. Liaquat MT, Jain S. Spontaneous Intracranial Hypotension. In: StatPearls [Internet]. Treasure Island : StatPearls Publishing; 2021
2. Pabaney AH, Mirza FA, Syed NA, Ahsan H. Spontaneous dural tear leading to intracranial hypotension and tonsillar herniation in Marfan syndrome: a case report. *BMC Neurol.* 2010;10:54. doi: 10.1186/1471-2377-10-54.
3. Nipatcharoen P, Tan SG. High thoracic/cervical epidural blood patch for spontaneous cerebrospinal fluid leak: a new challenge for anesthesiologists. *Anesth Analg.* 2011;113:1476-9. doi: 10.1213/ANE.0b013e31822e5665.
4. Ferrante E, Arpino I, Citterio A. Is it a rational choice to treat with lumbar epidural blood patch headache caused by spontaneous cervical CSF leak? *Cephalalgia.* 2006;26:1245-6. doi: 10.1111/j.1468-2982.2006.01179.x.
5. Cousins MJ, Brazier D, Cook R. Intracranial hypotension caused by cervical cerebrospinal fluid leak: treatment with epidural blood patch. *Anesth Analg.* 2004;98:1794-7, table of contents. doi: 10.1213/01.ane.0000117225.31416.36.
6. Kapoor SG, Ahmed S. Cervical epidural blood patch--A literature review. *Pain Med.* 2015;16:1897-904. doi: 10.1111/pme.12793.
7. Shim E, Lee JW, Lee E, Ahn JM, Kang Y, Kang HS. Fluoroscopically Guided Epidural Injections of the Cervical and Lumbar Spine. *Radiographics.* 2017;37:537-61. doi: 10.1148/rg.2017160043.