


High-resolution map for vegetation in the upper valleys and plateaus of Serra da Estrela (Portugal)

INÊS FERREIRA MENDES ^{1,2} CARLA MORA ^{1,2} SUSANA RODRÍGUEZ-ECHEVERRÍA ^{2,3} 

ABSTRACT – Mountain ecosystems such as Serra da Estrela, in the southwest of the Central Iberian Massif, are characterised by high ecological diversity shaped by climatic, geological, and anthropogenic factors. The vegetation of this region has been classified into distinct altitudinal belts, with the upper plateau (above 1600 m) hosting unique high-altitude scrubland and meadow communities of significant scientific interest. This study aimed to map and analyse the spatial distribution and temporal dynamics of vegetation in the upper valleys and plateaus of Serra da Estrela. High-resolution WorldView-2 satellite imagery from 2020 was used to produce a detailed classification of current vegetation, identifying eleven distinct classes. To assess vegetation changes over time, a comparative analysis was performed using a 1995 orthophotomap. The results reveal strong altitudinal zoning and spatial clustering of vegetation classes, with shrublands representing the dominant vegetation type (32.6%). Comparison with the 1995 data highlights an expansion of vegetated areas, particularly shrublands encroaching into former grassland zones. These findings demonstrate the value of high-resolution remote sensing for monitoring ecological changes in sensitive mountain environments and provide a crucial baseline for conservation and land management strategies in protected areas under pressure from land use shifts.

Keywords: Mountain vegetation; vegetation dynamics; remote sensing; WorldView-2.

RESUMO – MAPA DE ALTA RESOLUÇÃO DA VEGETAÇÃO DOS VALES E PLANALTOS SUPERIORES DA SERRA DA ESTRELA (PORTUGAL). Os ecossistemas montanhosos, como o da Serra da Estrela, no sudoeste do Maciço Ibérico Central, caracterizam-se por uma elevada diversidade ecológica moldada por fatores climáticos, geológicos e antrópicos. A vegetação desta região foi classificada em andares altitudinais distintas, sendo que o planalto superior (acima dos 1600 m) abriga comunidades singulares de matos e prados de altitude, de relevante interesse científico. Este estudo teve como objetivo mapear e analisar a distribuição espacial e a dinâmica temporal da vegetação nos vales e planaltos superiores da Serra da Estrela. Foi utilizada uma imagem de satélite WorldView-2 de alta resolução, referente ao ano de 2020, para produzir uma classificação detalhada da vegetação atual, identificando onze classes distintas. Para avaliar as alterações na vegetação ao longo do tempo, foi realizada uma análise comparativa com base num ortofotomapa de 1995. Os resultados revelam um acentuado zonamento altitudinal e agrupamentos espaciais das classes de vegetação, com os matos a representarem o tipo dominante (32,6%). A comparação com os dados de 1995 destaca uma expansão das áreas vegetadas, especialmente com matos a invadirem zonas anteriormente ocupadas por prados. Estes resultados demonstram a importância da deteção remota de alta resolução para a monitorização de alterações ecológicas em ambientes montanhosos sensíveis, oferecendo uma base essencial para estratégias de conservação e gestão territorial em áreas protegidas sob pressão das alterações no uso do solo.

Palavras-chave: Vegetação de Montanha; dinâmica da vegetação; deteção remota, WorldView-2.

HIGHLIGHTS

- High-resolution mapping reveals dynamic vegetation changes in Serra da Estrela.

Recebido: 04/04/2024. Aceite: 10/01/2025. Publicado: 01/10/2025.

¹ Centro de Estudos Geográficos, Instituto de Geografia e Ordenamento do Território, Universidade de Lisboa, Edifício IGOT, R. Branca Edmée Marques, 1600-276, Lisboa, Portugal. E-mail: inesmendes2@campus.ul.pt, carlamora@campus.ul.pt

² Laboratório Associado TERRA, Instituto de Geografia e Ordenamento do Território, Universidade de Lisboa, Lisboa, Portugal.

³ Centro de Ecologia Funcional, Departamento de Ciências da Vida, Universidade de Coimbra, Coimbra, Portugal. E-mail: susanare@ci.uc.pt

- Imagery classification effectively distinguishes eleven vegetation classes.
- Significant expansion of shrublands, linked to declining grazing.
- Notable forest reduction due to fires and land-use shifts over 25 years.
- Study aids conservation planning in Serra da Estrela's protected and sensitive areas.

1. INTRODUCTION

1.1. Overview

1.1.1. Background and context

Mountain ecosystems are globally recognised for their ecological significance, biodiversity richness, and sensitivity to environmental changes (Grabherr *et al.*, 2011; Körner, 2004). These high-altitude environments often host unique plant communities, shaped by steep ecological gradients, climatic variability, and long-term geomorphological processes (Viviroli *et al.*, 2007). Their vegetation patterns are not only important indicators of ecological health but also reflect the ongoing interactions between natural factors and human land use (Beniston, 2003). The patterns observed here mirror broader trends in high-altitude environments worldwide, where climatic gradients, isolation, and traditional land-use practices have shaped rich but vulnerable plant communities.

Vegetation mapping in mountain areas has become increasingly relevant in the context of climate change and land-use transformations, which are causing notable shifts in species composition, distribution, and structure (Pauli *et al.*, 2012; Tasser *et al.*, 2005). Accurate and up-to-date vegetation cartography is essential for understanding these dynamics and for guiding conservation and sustainable land management policies in protected mountain landscapes.

1.1.2. Remote Sensing in Vegetation Mapping

The need to improve the quality and detail of cartography has led to the application of Remote sensing (RS) methods that revolutionised the study of vegetation, particularly in morphologically complex and inaccessible areas such as mountainous regions. The use of satellite imagery provides access to a wide range of multispectral data, offering extensive temporal and spatial coverage at a relatively low cost, especially with open-source options available (Monteiro-Henriques *et al.*, 2016; Xie *et al.*, 2008). The high-resolution and repeatability of RS data make it a practical solution for studying the dynamic and diverse environments of mountain ecosystems. Furthermore, RS technologies allow for the identification of species and communities, detection of changes in vegetation distribution patterns, providing insights into the dynamics of these communities and the effects of external factors such as climate change and human activities (Wang *et al.*, 2015). However, applying RS in mountainous regions presents specific challenges due to the complexity of the landscape, including steep topography, rapid and significant environmental changes, and atmospheric and topographic conditions that can affect data accuracy (Gartzia *et al.*, 2013). Despite these challenges, RS remains an invaluable tool for environmental monitoring and natural resource management in these difficult-to-access areas. In this specific work, a multispectral image from August 2020, provided by a satellite (Worldview 2), with 2m resolution, was mainly used to classify the vegetation communities on the plateau. Due to the flowering period of a specific community, a Sentinel-2 image was also used to differentiate it from other vegetation.

1.1.3. Objectives of the Current Study

Despite the existing studies, a high-resolution mapping of the vegetation in this area has not been undertaken until now. The work presented here complements previous studies by providing a detailed cartographic analysis of the most significant vegetation communities on the plateau of Serra da Estrela. In addition, this study also aims to assess the vegetation dynamics by comparing the newly produced map with previous photographic records, such as ortho-photomaps from 1995. This analysis

will provide insights into the temporal changes and shifts in vegetation cover, allowing for a better understanding of how these ecosystems have evolved over time. This cartographic work is fundamental scientific knowledge that supports better land management practices to protect biodiversity and mitigate potential disturbances to the habitats in this area.

1.2. Study area

1.2.1. Geographical Context

The Serra da Estrela (40°20'N, 7°35'W) is part of the Central Mountain Range. It belongs to the Hesperic Massif, forming the westernmost and highest part of a mountain alignment extending SW-NE, composed of elongated plateaus-oriented WSW-ESE (fig.1). This region stretches from Guarda to Serra da Lousã, covering approximately 115km with an average width of 25km (Daveau, 1969). It is internationally recognized for its geological significance and is part of the Unesco Geoparks Network (Estrela Geopark). Additionally, it hosts the Serra da Estrela Natural Park, the largest protected area in Portugal, which shelters endemic vegetation and wildlife refuges of national importance (*Decreto-Lei n.º 557/76*). The specific study area within this context (fig.1 at the bottom left) focuses on the Upper Plateau, from Vale do Rossim to Torre, including the headwaters of the Zêzere, Alforja, Loriga, Caniça, and Alva rivers. This area is of significant scientific, environmental, landscape, and cultural interest and is included in the Natura 2000 network, Ramsar Convention, and the biogenetic reserve of the Serra da Estrela Natural Park (Castro *et al.*, 2019).




Fig. 1 – Framing of Estrela Geopark, Serra da Estrela Natural Park and the Study area.

Fig. 1 – Enquadramento do Geopark Estrela, do Parque Natural da Serra da Estrela e da área de estudo.

Source: Authors' elaboration

1.2.2. Geological and Topographic Features

Serra da Estrela is the highest mountain range in mainland Portugal, with the highest elevation in the southwest, reaching 1993m at the Planalto da Torre (Vieira, 2004). The mountain is bounded by fault scarps to the east and west, and river incisions shape its current topography following the uplift of the mountain. The landscape is also heavily influenced by the nature of its rocks, mainly granite and schist, as well as by the glacial activity during the Last Glacial Maximum in the Upper Pleistocene (Ferreira *et al.*, 2001; Ferreira & Vieira, 1999; Vieira, 2004). The higher plateau, above 1600m, is divided by the Zêzere and Alforfa valleys into two main sectors: the Western and Eastern Plateaus. The study area is within the Western Plateau, characterized by distinct relief subunits, such as Planalto das Penhas Douradas (1500-1550m), Planalto do Curral do Martins (1650-1760m), and Planalto da Torre (above 1800m), with the Curral do Martins and Torre plateaus being referred to as the Superior Plateau (Vieira, 2004).

1.2.3. Climate Characteristics

The climate of Serra da Estrela is distinctive and varies significantly with altitude, topography, and exposure to prevailing winds. According to the Köppen classification the the southeast side of Serra da Estrela Mountain is characterized by a hot-summer Mediterranean (Csa) type, whereas the northwest side features a mild-summer Mediterranean (Csb) climate (Santos *et al.*, 2024). The mountain's altitude, topography, and orientation play significant roles in modifying the local climates, with increased altitude leading to lower annual average temperatures, estimated to decrease by 0.6°C per 100m of elevation (Mora, 2006). The study area, specifically the Upper Plateau, has an estimated average annual temperature of 4-6°C, with the coldest month being January and the warmest month being July. Precipitation is marked by abundant rainfall in winter and dry summers. The wet season is from October to May, with a mean annual precipitation of ca. 2500mm in the summit, while the plateaus show more than 2000mm. The western side of the mountain presents a larger number of days with rainfall, but a slightly lower total amount than the eastern part, which in turn shows a smaller number of days with rain (Daveau *et al.*, 1977). The western slopes receive more precipitation due to orographic lifting, contrasting windward and leeward areas (Vieira & Mora, 1998). Frequent fogs on the western slope are responsible for hidden precipitation that increases water availability. Snowfall is a significant feature in the higher altitudes, though it varies greatly in frequency and duration. The presence of snow plays a crucial role in protecting vegetation from extreme cold and wind, contributing to soil and plant temperature regulation, and supporting plant growth during the melting period (Pauli *et al.*, 2012; Wang *et al.*, 2015).

1.2.4. High-Altitude Vegetation in Serra da Estrela

The high-altitude plateau of Serra da Estrela, precisely above 1600m, presents a distinct type of vegetation compared to lower altitudes. This area is predominantly characterized by shrublands and high-altitude grasslands typical of mountain environments (Jansen, 2002). The vegetation here is shaped by extreme climatic conditions, including thermal extremes, snow accumulation for longer periods, and wind, alongside historical and ongoing anthropogenic influences such as pastoral activities, intentional burning, and more recently, tourism (Jansen, 2011).

The current vegetation in Serra da Estrela is also the result of significant climatic changes over time, with historical evidence indicating the presence of ten distinct climatic and edaphic-forest types, many of which have been altered or destroyed (van der Knaap & van Leeuwen, 1994). Fire, both natural and anthropogenic, has played a crucial role in shaping the vegetation, often acting as both a destructive force and a process of pasture renewal, though its increasing frequency in recent decades poses a significant threat to local ecosystems (Barrico *et al.*, 2019; Connor *et al.*, 2012). Tourism has further impacted the landscape by increasing ecosystem pressures through recreational activities (Fidalgo, 1994; Jansen, 2011; Soncco, 2020). The unique geographical and climatic conditions of Serra da Estrela have made its flora a scientific interest since the 18th century, with significant botanical expeditions contributing to our understanding of this isolated mountain's vegetation (Machado *et al.*, 2007).

The region hosts over 900 taxa from more than 90 botanical families, including several endemic species and a variety of floristic elements such as Mediterranean, Temperate European, Boreo-Alpine,

and Arctic-Alpine types, all of which have been significantly altered by human activity (Jansen, 2002; Meireles & Pinto-Gomes, 2012). The richness of its flora is attributed to factors like climate variability, geological diversity, proximity to Africa and the Atlantic, and historical influences, particularly the Quaternary glaciations (Jansen, 2002; 2011).

1.2.5. Historical and Botanical Significance of the Region



The unique characteristics of this vegetation have long attracted the interest of the scientific community, resulting in a significant amount of research that highlights the diversity and specificity of the flora in this region (Henriques, 1883; Meireles, 2010; Meireles, *et al.*, 2008; Pinto da Silva & Teles, 1980). In 1994, J.P. Fidalgo mapped the vegetation above 1600 meters in Serra da Estrela, using a grid system to document the distribution of species prioritized for conservation, providing a detailed inventory of species (Fidalgo, 1994). The Institute for Nature Conservation and Forests [ICNF] produced later a different vegetation map, focusing on natural habitats of community interest, as listed in the Habitats Directive. This mapping predominantly identified heath and shrub habitats in temperate zones and flora of ecological interest, although it differed from Fidalgo's classification ([ICNF], 2019).

1.2.6. Vegetation Types and Habitats

The vegetation of Serra da Estrela is diverse, with significant altitudinal variation, influencing the distribution of plant communities. The area of interest, the Upper Plateau, is predominantly characterized by high-altitude shrublands and grasslands, which dominate above 1600m. These habitats are of considerable biodiversity importance and are included in conservation initiatives such as Natura 2000 (table I), Ramsar, and the biogenetic reserve of the Serra da Estrela Natural Park. Shrublands in the upper areas are dominated by species such as *Juniperus communis* var *saxatilis* Pall., *Cytisus oromediterraneus* L. and *Echinopartum ibericum* Rivas Mart., Sánchez Mata & Sancho. Between 1400 and 1700m of altitude, dominant species include other leguminous shrubs such as *Cytisus stratus* L., *C. scoparius* L., *G. florida* L. or *G. cinerascens* L., and the heaths *Erica arborea* L. or *E. australis* L. (Jansen, 2002). Other smaller shrubs such as *Calluna vulgaris* L. or *Halimium lasianthum* subsp *alyssoides* (Lam.) Greuter. can be found associated with the main shrubs. Grasslands, maintained by grazing, are mostly dominated by *Nardus stricta* L., although they can harbor a significant number of endemic and rare species such as *Festuca henriquesii* Hack, *Campanula herminii* Hoffmans & Link, *Armeria sampaioi* (Bernis) Nieto Fel., *Gentiana lutea* L. or *Fritillaria nervosa* subsp *nervosa* Willd.

Table I – Main classified communities (in terms of occupied area), respective examples in the field and Natura 2000 habitat (cont.). Colour figure available online.





Quadro I – Principais comunidades classificadas (em termos de área ocupada) e respetivos exemplos no terreno e habitat Natura 2000 (cont.). Figura a cores disponível online.

Field Photographs	Plant Communities	Main characteristic species	Natural and Semi-natural habitats (Natura 2000 level)
	Matgrass Grasslands	<i>Nardus stricta</i> L.; <i>Festuca henriquesii</i> ; <i>Festuca rubra</i> L.; <i>Juncus squarrosus</i> L.	6230- Species-rich <i>Nardus</i> grasslands, on silicious substrates in mountain areas (and submountain areas in Continental Europe)
	Shrubland	<i>Juniperus communis</i> var. <i>saxatilis</i>	4060- Alpine and Boreal heaths

Source: Authors' elaboration

Table I – Main classified communities (in terms of occupied area), respective examples in the field and Natura 2000 habitat. Colour figure available online.

Quadro I – Principais comunidades classificadas (em termos de área ocupada) e respetivos exemplos no terreno e habitat Natura 2000. Figura a cores disponível online.

Field Photographs	Plant Communities	Main characteristic species	Natural and Semi-natural habitats (Natura 2000 level)
	Alpine shrubland	<i>Cytisus oromediterraneus</i> L.	5120- Mountain <i>Cytisus purgans</i> formations
	Heaths	<i>Erica australis</i> subsp. <i>aragonensis</i> (Willk.) Count; <i>Erica arborea</i> L.; <i>Calluna vulgaris</i> L.; <i>Halimium alyssoides</i> (Lam.) Spach; <i>Ulex minor</i> ; <i>Ulex europaeus</i> subs. <i>latebracteatus</i> .	4030- European dry heaths
			
	Brooms	<i>Genista florida</i> L.; <i>Genista cinerascens</i> L.; <i>Cytisus striatus</i> L.; <i>Cytisus scoparius</i> L.; <i>Cytisus multiflorus</i> L.	-

Source: Authors' elaboration

2. METHODOLOGY

The methodology of this investigation was structured in three main phases (fig. 2). Phase I involved a literature review, selection and analysis of the study area, fieldwork, and the selection and acquisition of images for vegetation classification and comparison. Phase II focused on image pre-processing, followed by the classification of vegetation communities and creation of the final maps. Finally, in Phase III (section IV Discussion), vegetation dynamics over the past decades were interpreted by comparing the 1995 ortho-photomap with the 2020 satellite imagery, allowing for the identification of significant changes in the vegetation communities over time.

2.1. Phase I

2.1.1. Fieldwork

Fieldwork was fundamental to the classification process, involving three campaigns for data collection and validation (fig. 3). The initial campaign, conducted in September 2021, focused on collecting ground truth points using differential GPS (D-GPS) to establish training areas for image classification.

Despite challenges like limited network coverage in higher elevation areas, supplementary methods such as hand-held GPS devices and photographic documentation facilitated the accurate identification of vegetation types. Subsequent field campaigns in April and May 2022 were dedicated to collect validation points to ensure the accuracy of the vegetation maps.




Fig. 2 – Phases of research work.
 Fig. 2 – Fases do trabalho de investigação.
 Source: Authors' elaboration

2.1.2. Data Collection

For the vegetation classification, two satellite images and one ortophotomap were used. For the 2020 cartography of the plateau, high-resolution satellite imagery was used from WorldView-2 (WV2).

WV2 offers high spatial resolution with 2.0 meters in eight multispectral bands, including standard RGB and NIR bands, as well as additional bands like coastal blue, yellow, red-edge, and a second near-infrared (NIR2) band (table II), which are particularly useful for vegetation analysis (Hung & Treitz, 2020; Rapinel *et al.*, 2014). The WV2 image used in this study was captured on August 1, 2020, under optimal conditions—clear skies, minimal shadow, and no cloud cover. Due to the inability of this image to distinguish all the desired vegetation classes found in the field, and solve spectral overlapping between them, an S2 image of the flowering season was used.

To understand the recent changes in vegetation distribution on the Serra da Estrela Plateau a comparison from 1995 and 2020 was made. This comparison was essential to understand the dynamics of vegetation communities over 25 years. This study utilized orthoimages from 1995, obtained from the Directorate General for Territory. The image, with a scale of 1:25 000 and a resolution of 1 meter, lacked sufficient spectral information for automated classification (table II).

2.2. Phase II

2.2.1. Image pre-processing

The WV2 image pre-processing included radiometric and atmospheric corrections to enhance the quality of the WV2 images. Radiometric calibration adjusted the images from radiance to reflectance and was performed using the Radiometric Calibration tool in the ENVI software (Harris Geospatial, 2023). The atmospheric correction, was applied using the FLAASH tool in the same software, using the image acquisition information (Latitude, Longitude, date and time of the flight and sensor type), was also defined the atmospheric model to Mid-Latitude Summer (MLS), a rural aerosol model and 40km for the initial visibility, considering a clear sky at the acquisition time (Harris Geospatial, 2023). This improved the accuracy by accounting for atmospheric conditions at the time




Fig. 3 – Survey of the sampling points of the plant communities and validation, using GPS, throughout the study area and in detail in the Nave de Santo António. Photo of the fieldwork. Colour figure available online.

Fig. 3 – Levantamento dos pontos de amostragem das comunidades vegetais e validação, através de GPS, em toda a área de estudo e em pormenor na Nave de Santo António. Fotografias de campo. Figura a cores disponível online.

Source: Inês Mendes (2021)

Table II – Sensors used and their characteristics (images used, camera, bands with central wavelength and resolution, represented at the same scale).

Quadro II – Sensores utilizados e respetivas características (imagens utilizadas, câmara, bandas e central wavelength, altura do voo e resolução, representadas à mesma escala).

	Satellites		Orthophotomap
	WorldView-2	Sentinel-2	
Camera	WorldView-110	Sentinel (L2A image)	
Acquisition date	01/08/2020	30/05/2020	1995
Bands	Coastal (400-450 nm) Blue (450-510 nm) Green (510-580 nm) Yellow (585-625 nm) Red (630-690 nm) Red Edge (705-745 nm) Near-Infrared 1 (770-895 nm) Near-Infrared 2 (860-1,040 nm)	Coastal Aerosol (443 nm) Blue (490 nm) Green (560 nm) Red (665 nm) Red Edge 1 (705 nm) Red Edge 2 (740 nm) Red Edge 3 (783 nm) Near-Infrared (NIR) (842 nm) Narrow Near-Infrared (865 nm) Water Vapor (945 nm) SWIR-Cirrus (1,375 nm) Short-Wave Infrared 1 (SWIR 1) (1,610 nm) Short-Wave Infrared 2 (SWIR 2) (2,190 nm)	False color
Resolution	Panchromatic ~ 0.5m Multispectral ~ 2.0m	10m	1m
Source	ESA	Copernicus EO browser	https://www.dgterritorio.gov.pt/

Source: Authors' elaboration

of image capture (Adam *et al.*, 2017; Brunn *et al.*, 2003). There was no need to perform orthorectification since the image is already extracted with this correction (Rapinel, *et al.* 2014). For the Sentinel-2 L2A the pre-processing process was not necessary, as this image is already fully corrected – including orthorectification, and radiometric, atmospheric, and topographic corrections.

Orthophotomaps were downloaded as individual images, requiring georeferencing, orthorectification, colour correction and harmonization to be suitable for subsequent classification. The entire process was conducted in a GIS environment using ArcGIS Pro software (Esri, 2023). The first step involved georeferencing each individual image using the spline transformation method, based on known coordinate points within the image. To produce the final mosaic, the images were combined using the *Mosaic Dataset* tool with default parameters. To enhance the visual quality of the final mosaic–by homogenising colour, brightness, and contrast– histogram manipulation and colour correction was performed using the *Dodging* method for colour balance and the *first-order* method for colour surface type. These steps were critical to ensure the extraction of high-quality data, following the guidelines of Brunn *et al.* (2003).

2.2.2. Vegetation Classification Process

For the WV2 and S2 image classification process, a supervised approach (Rapinel *et al.*, 2014) was employed, utilizing Regions of Interest (ROIs) based on field data and photointerpretation, using all the bands available to better distinguish between classes (Hung *et al.* 2020). To determine the number of classes needed, a preliminary unsupervised classification was conducted, providing an estimate of the optimal number of vegetation classes for detailed mapping. This step facilitated the *pixel based* supervised classification, which was considered the most suitable method for this type of vegetation analysis. Several classifiers were tested, including parametric methods like Maximum Likelihood, which achieved the best overall accuracy. The final optimal number of classes was achieved after various attempts and validation on the field (see section 2.3).

The orthophotomap was processed through manual photointerpretation and classification of the final mosaic. Vegetation classes were harmonized considering the main vegetation units, resulting in five broad categories: Forest, Shrubs, Matgrass grasslands, Spare Grasslands, and "Other Classes".

2.2.3. Validation

Validation was carried out using two approaches: statistical and in-situ. No validation was performed for the orthophotomap, as the classification was manual and to the year 1995.

Statistical accuracy assessment was conducted through the generation of an error matrix. To produce the error matrix, new ROIs were created for each class used in the final classification, based on field samples. Using the post-processing tool in ENVI software, the matrix was generated (table III) with the *Confusion Matrix Using Ground Truth ROIs* tool.

In-situ validation was carried out in fieldwork and became crucial to understand and correct significant spectral confusion that persisted among vegetation classes in the WV2 classification.

After the validation process the initial classification was refined, supported on knowledge of the study area. Certain Shrubland classes, such as *Cytisus*, *Genista* and Heaths were merged due to overlapping spectral characteristics (originating the 'Brooms and Heaths' class), especially outside the flowering season, and their dense distribution in mixed vegetation communities. These adjustments helped reduce classification errors, improving the overall map accuracy to 94% (table III).

To address the necessity of merging classes, the Sentinel-2 image classification was integrated, using all the available bands and following the same methodological procedure as the WV2. This last was particularly useful for distinguishing certain species like Heaths (belonging to the class 'Brooms and Heaths' in the WV2 classification) during their flowering period, illustrating the importance of selecting temporally appropriate imagery (Anderson *et al.*, 1993; Cingolani, 2004). Despite challenges, including the need to merge similar classes and adjust for spectral variations within the same species, the methodology enabled a detailed and accurate mapping of vegetation communities on the plateau. The process underscored the critical role of combining field data, advanced satellite imagery, and sophisticated classification techniques in ecological mapping.

Table III – Validation of the classification carried out for the Plateau (WorldView 2 image).

Quadro III – Validação da classificação efectuada para o Planalto (imagem WorldView 2).

Classification classes	Producer Accuracy (%)	User Accuracy (%)
Evergreen Forest	78.35	60.32
Deciduous Forest	96.27	91.49
Brooms and Heaths	81.25	89.97
- Heaths*	83.13	79.50
Rockroses	90.91	64.52
Heathers	97.55	80.78
Juniper	84.00	61.76
Matgrass grasslands	88.14	97.17
Sparse Matgrass	83.54	42.58
Rushes	100.0	93.33
Ferns	83.14	85.22
Bare soil	50.50	42.86
Bare Rock	73.17	71.31
Water Bodies	99.43	99.99
Shadow	99.04	56.28
Road	58.88	93.93
Built Area	87.31	94.55
Agricultural area	97.60	89.56
Overall Accuracy = 93.7%		
Kappa Coefficient = 0.85		

*Heaths classified with the Sentinel 2 image

Source: Authors' elaboration based on WorldView 2 image

3. RESULTS

3.1. Current Characterization of Vegetation Communities

Bare rock accounts for 32% of the area, being the second most represented class which reflects the high-altitude terrain with minimal vegetation cover. Rock outcrops are often covered with lichens and mosses. Other non-vegetation classes include areas of bare soil, where sparse grasses grow among the granite sands, and a minimal presence of built structures, mainly related to tourism activities at the mountain summit and the village of Sabugueiro in the northwest of the classified area. Water bodies on the plateau, such as natural, semi-natural, and artificial lagoons, are mapped. Large flooded areas, like peat bogs, which contain other plant communities, were not included in the mapping.

Our study identified eleven vegetation classes within the mapped area (fig. 4). These classes range from forests at lower altitudes to communities of Juniper, Heather, and matgrass at higher elevations. Forests are located at lower altitudes (1530m for evergreen forests and 1600m for deciduous forests), and a gradual transition to Alpine Shrubland and Grassland communities at higher altitudes.

Shrublands and Alpine Shrublands are divided into five classes: Brooms (*Cytisus* and *Genista* spp), Heaths (*Erica* spp), Heathers (*Calluna vulgaris*), Rockroses (*Halimium lasianthum subsp alyssoides*) and Junipers (*J. communis* var. *saxatilis*). Brooms and Heaths together represent the largest vegetation type representing 65.9% of the total vegetation cover. *Cytisus* and *Genista* species, classified together as Brooms, are widespread across the plateau, occupying most of the vegetated territory, particularly along valleys, slopes, and areas around Vale do Conde.

The heath class was classified separately due to a different image resolution (10m compared to 2m for the rest of the map), and its distribution often overlaps with other classes on the ground. Heaths

are primarily found between 1600 and 1700m, while other shrub species (also belonging to the heathland community) like rockroses and heather are associated with moist areas of the plateau, such as Nave de Santo António, Vale da Candeeira, and Vale do Conde. Junipers represents the upper altitudinal range of vegetation distribution and is strongly represented at altitudes between 1700 and 1900m, particularly along the road connecting Torre to Lagoa Comprida and on the area below Torre.




Fig. 4 – Distribution of the communities under study for the Serra da Estrela Plateau and their mapped percentage.

*Different resolution (S2 derived). Colour figure available online.

Fig. 4 – Distribuição das comunidades em estudo para o Planalto da Serra da Estrela e respetiva percentagem cartografada. *Resolução Diferente (S2 derived). Figura a cores disponível online.

Source: Authors' elaboration based on WorldView 2

Grasslands, either densely covered by matgrass or sparsely covered by other grasses such as *Agrostis truncatula* Parl, are mainly found in depressed areas, mostly above 1600m, and in surrounding areas like Nave de Santo António, Lagoa Comprida, and Vale do Conde. These communities are often spatially associated with other species such as Rockroses, Heaths, Heath rushes (*Juncus squarrosus* L.) and sometimes Gorses (*Ulex* spp). Sparse grasslands, which occur throughout the upper plateau, are interspersed with bare soil.

Rushes and Ferns were mapped due to their distinct spectral signatures and significant presence in the area. Rushes are less visible at this scale, mainly in grasslands or areas with high soil moisture. Ferns, which are more prominent, originate from two distinct communities: those in recently burned areas dominated by *Pteridium aquilinum* and those in moist, sheltered locations along valley floors dominated by *Polystichum setiferum*. Their main expression is between 1300 and 1600m, with sporadic occurrences above 1600m.

Echinopartum ibericum, an endemic spiny legume shrub locally known as “Caldoneira” that appears associated with junipers and high-altitude brooms was not classified due to its residual distribution and the resolution used in the classified image.

3.2. Comparison of Vegetation Dynamics (1995-2020)

Comparing the orthophotomap obtained for 1995 and the high-resolution 2020 map (fig. 5), we observed a notable decrease in forested areas, particularly in the northern region around the village of Sabugueiro and valleys, which had dense tree cover in 1995 but by 2020, these zones exhibited a visible increase in scrubland and sparse scrubland, suggesting a shift in vegetation cover.




Fig. 5 – Mapping of the plateau communities for 1995 and 2020.
Fig. 5 – Cartografia das comunidades do planalto para 1995 e 2020.

Source: Authors' elaboration based on WorldView 2

Similarly, Fern-dominated areas, classified under "Other Communities," have diminished in size and are now mixed with shrubs and patches of forest. Grasslands, previously widespread across the central and southern sectors—particularly near Alto da Torre and Lagoa Comprida – showed a marked

decrease in extent. Agricultural land has also been reduced, especially near Sabugueiro, highlighting a shift in land use. The shrublands showed a mixed pattern, with some areas like the Conde Valley exhibiting reduced shrub density, while other regions, such as near Lagoa Comprida, experienced an increase in shrub density over time. Built-up areas and the road network remained largely unchanged across both maps.

4. DISCUSSION

This work presents a detailed study of the spatial distribution and temporal dynamics of the vegetation in Serra da Estrela upper valleys and plateaus. The mapping of vegetation communities contributes to the delimitation of habitats with relevant natural value, such as those limited by the Natura 2000 Network Sector Plan, supporting decision-making and conservation planning. Also, the vegetation in these extreme environments serves as an indicator of the changes taking place at the level of the ecosystem in which they are located and can be vulnerable to changes and disturbances.

Methodologically, the use of very high-resolution, multispectral satellite imagery – with eight distinctive bands – for the main supervised classification proved to be crucial for the detailed mapping of the plateau, achieving an overall accuracy of 94%. This distinguishes the study from others relying on lower-resolution data or field surveys alone, emphasizing the value of Earth Observation in environmental monitoring.

Nevertheless, the classification faced challenges, particularly in spectral separation of certain species and communities, and the exclusion of some less represented classes due to the 2m pixel resolution. It was impossible to separate *Cytisus*, *Genista* and *Erica* due to their proximity on the ground and spectral similarity. However, despite *Cytisus* and *Genista* species being physiognomically similar, they form communities with different characteristics and are separated in the territory, with *Cytisus oromediterraneus* dominating at higher altitudes.

Among the limitations of this study is the use of an WV2 imagery from a single season, which may reduce the accuracy of classification for species with seasonal variability, and the difficulty in mapping small or fragmented vegetation types that fall below the 2m pixel resolution. Improvements could be achieved with high-resolution imagery from different seasons, enabling more detailed classification and the distinction of geographically significant species. Field-based validation across seasons and more extensive ground-truth datasets could also refine classification accuracy and support long-term ecological monitoring.

Despite those limitations, the analysis provided a comprehensive understanding of vegetation patterns across the study area, revealing clear spatial relationships between different plant communities. These patterns are strongly influenced by altitude, with distinct variations in vegetation distribution that reflect gradients along altitudinal zones.

A noticeable decrease in forest cover, likely linked to recurring forest fires, such as the significant event in 2017, marks one of the key vegetation changes observed over the study period. However, monitoring forest areas is of special interest to detect potential shifts in altitude, particularly in the context of climate change, which could drive forest expansion above 1600m in protected areas (Grabherr *et al.*, 1994; Pauli *et al.*, 2003).

The most significant change was observed in the class Matgrass grasslands, which have substantially declined, with these areas being increasingly overtaken by shrubs. This shift may be linked to the decline in pastoral activities, as grazing and controlled burns has historically maintained these grasslands and limit the shrub expansion. If we analyse the number of livestock by geographic location (Instituto Nacional de Estatística [INE], 2022) within the municipalities that the study area is part of (primarily Seia, followed by Manteigas), we can observe a decrease in the number of sheep indicating a reduction on pressure from grazing activities. This reduction has likely contributed to the expansion of shrublands, as grazing is responsible for maintaining Matgrass grasslands by controlling shrub growth (Rudley *et al.*, 2023; Sanz-Elorza *et al.*, 2003). The loss of these practices is now driving to passive ecological succession toward more homogeneous and woody vegetation, with implications in biodiversity, affect soil infiltration and water retention, and increase flammability, potentially amplifying the frequency and intensity of wildfires in the region.

Along with the decline in grazing, climate change may be also promoting shrub encroachment as it has been shown for other mountains. Indeed, our results align with the recorded expansion of *C. oromediterraneus* and *J. communis* in the upper belt of the Iberian Central Massif and the Moncayo

mountain, where shrub encroachment into grasslands has been noted, influenced by factors such as rising temperatures, changes in land use, and reduced grazing (Gómez-García *et al.*, 2023; Sanz-Elorza *et al.*, 2003).

5. CONCLUSION

The use of high-resolution satellite imagery and advanced GIS techniques, supported the successful mapping of vegetation of the Serra da Estrela Upper Plateau, providing a detailed representation of the vegetated area. This mapping reveals a diverse and complex vegetation landscape, shaped by natural and anthropogenic factors. This study also reveals alterations in the vegetation over a short period (1995-2020), marked by a reduction in forested and agricultural areas and an increase in shrubland. Factors such as large fires and shifts in land use, particularly the decline in traditional pastoral practices and the reduced use of fire for pasture maintenance, may have also contributed to the spread of shrubland into areas previously dominated by grasslands. These findings are consistent with similar trends observed in other Mediterranean and montane environments, where land abandonment have driven shrub invasion and a reduction in traditional open habitats (Brandt, *et al.*, 2013; Gómez-García *et al.*, 2023; Pauli *et al.*, 2003; Sanz-Elorza *et al.*, 2003).

Overall, this research contributes with valuable insights into the ecological structure and dynamics of this protected area, useful for future land management and conservation strategies aimed at preserving its rich natural heritage. The detailed knowledge of the distribution of the main habitats is the first step towards adopting appropriate management measures, making it possible to preserve the open landscape and this vegetation, which is of scientific and environmental interest.

ACKNOWLEDGEMENTS

This work was funded by the Programmatic Budget of the Centro de Estudios Geográficos (UIDP/00295/2020). Special thanks are to the Estrela Geopark for the assistance during fieldwork.

ORCID ID

Inês Ferreira Mendes  <https://orcid.org/0000-0001-5345-0097>

Carla Mora  <https://orcid.org/0000-0002-0843-3658>

Susana Rodríguez-Echeverría  <https://orcid.org/0000-0002-2058-3229>

AUTHOR'S CONTRIBUTION

Inês Ferreira Mendes: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Resources, Data curation, Writing – original draft preparation, Writing – review and editing, Visualization. **Carla Mora:** Conceptualization, Methodology, Validation, Investigation, Resources, Data curation, Writing – review and editing, Visualization, Supervision, Project administration, Funding acquisition. **Susana Rodríguez-Echeverría:** Conceptualization, Validation, Writing – review and editing, Supervision.

REFERENCES

- Adam, E., Mureriwa, N., & Newete, S. (2017). Mapping *Prosopis glandulosa* (mesquite) in the semi-arid environment of South Africa using high-resolution WorldView-2 imagery and machine learning classifiers. *Journal of Arid Environments*, 145, 43-51. <https://doi.org/10.1016/j.jaridenv.2017.05.001>
- Anderson, G.L., Everitt, J.H., Richardson, A.J. and Escobar, D.E., (1993). Using satellite data to map false broomweed (*Ericameria austrotexana*) infestations on south. Texas rangelands. *Weed Technology*, 7(4), 865-871. <https://doi.org/10.1017/S0890037X00037908>

- Barrico, L., Freitas, H., & Rodríguez-Echeverría, S. (2019). *Caracterização demográfica e cartográfica das aldeias de montanha: Potenciais riscos ambientais e socioeconómicos*. [Demographic and cartographic characterisation of mountain villages: Potential environmental and socio-economic risks]. Universidade de Coimbra.
- Beniston, M. (2003). Climatic change in mountain regions: A review of possible impacts. *Climatic Change*, 59(1-2), 5–31. <https://doi.org/10.1023/A:1024458411589>
- Brandt, J. S., Haynes, M. A., Kuemmerle, T., Waller, D. M., & Radeloff, V. C. (2013). Regime shift on the roof of the world: Alpine meadows converting to shrublands in the southern Himalayas. *Biological Conservation*, 158, 116–127. <https://doi.org/10.1016/j.biocon.2012.07.026>
- Brunn, A., Fisher, C., Dittmann, C., & Richter, R. (2003). Quality assessment, atmospheric and geometric correction of airborne hyperspectral HyMap data. In *3rd EARSeL Workshop on Imaging Spectroscopy* (pp. 13–16). Technical University of Clausthal. <https://elib.dlr.de/455/>
- Castro, E., Gomes, H., Loureiro, F., Fernandes, M., & Patrocínio, F. (2019). Estrela Geopark: Uma Estratégia de Desenvolvimento Territorial [Estrela Geopark: A Territorial Development Strategy]. *Praça Velha*, 39(1), 271–288. <http://hdl.handle.net/10400.6/10274>
- Cingolani, A. M., Renison, D., Zak, M. R., & Cabido, M. R. (2004). Mapping vegetation in a heterogeneous mountain rangeland using landsat data: an alternative method to define and classify land-cover units. *Remote Sensing of Environment*, 92, 84–97. <https://doi.org/10.1016/j.rse.2004.05.008>
- Connor, S. E., Araújo, J., van der Knaap, W. O., & van Leeuwen, J. F. N. (2012). A long-term perspective on biomass burning in the Serra da Estrela, Portugal. *Quaternary Science Reviews*, 55, 114–124. <https://doi.org/10.1016/j.quascirev.2012.08.007>
- Daveau, S. (1969). Structure et relief de la Serra da Estrela [Structure and relief of the Serra da Estrela]. *Finisterra – Revista Portuguesa de Geografia*, IV(8) 5–32. <https://doi.org/10.18055/Finis2482>
- Daveau, S., Coelho, C., Costa, V., & Carvalho, L. (1977). *Répartition et rythme des précipitations au Portugal* [Distribution and rate of rainfall in Portugal]. *Memórias do Centro de Estudos Geográficos*, 3. Centro de Estudos Geográficos, Universidade de Lisboa.
- Esri. (2023). ArcGIS Pro [Computer software]. Environmental Systems Research Institute. <https://www.esri.com>
- Ferreira, A. B., Alcoforado, M. J., Vieira, G. T., Mora, C., & Jansen, J. (2001). *Landscape analysis and classification methodologies: The example of the Estrela Project*. *Finisterra – Revista Portuguesa de Geografia*, XXXVI(72), 157–178. <https://doi.org/10.18055/Finis1632>
- Ferreira, N., & Vieira, G. (1999). *Geological and geomorphological guide of the Serra da Estrela Natural Park: Sites of geological and geomorphological interest*. Instituto da Conservação da Natureza
- Fidalgo, J. P. P. (1994). A distribuição da flora vascular na área do Parque Natural da Serra da Estrela acima dos 1600 metros de altitude [The distribution of vascular flora in the Serra da Estrela Natural Park area above 1600 metres above sea level]. In *II Seminário Técnico: Conservação da Natureza na Serra da Estrela. Comunicações*. Instituto da Conservação da Natureza.
- Gartzia, M., Alados, C. L., Pérez-Cabello, F., & Bueno, C. G. (2013). Improving the accuracy of vegetation classifications in mountainous areas. *Mountain Research and Development*, 33(1), 63–74. <http://dx.doi.org/10.1659/MRD-JOURNAL-D-12-00011.1>
- Gómez-García, D., Aguirre de Juana, Á. J., Jiménez Sánchez, R., & Manrique Magallón, C. (2023). Shrub encroachment in Mediterranean mountain grasslands: Rate and consequences on plant diversity and forage availability. *Journal of Vegetation Science*, 34, e13174. <https://doi.org/10.1111/jvs.13174>
- Grabherr, G., Gottfried, M., & Pauli, H. (1994). Climate effects on mountain plants. *Nature*, 369, 448. <https://doi.org/10.1038/369448a0>
- Grabherr, G., Gottfried, M., & Pauli, H. (2011). Global Change Effects on Alpine Plant Diversity. In F. E. Zachos & J. C. Habel (Eds.), *Biodiversity Hotspots* (pp. 149–163). Springer.
- Harris Geospatial (2023). *ENVI* [Computer software]. NV5 Global.
- Henriques, J. A. (1883). *Expedição científica à Serra da Estrela em 1881: Secção de botânica* [Scientific expedition to Serra da Estrela in 1881: Botany section]. Sociedade de Geografia de Lisboa.
- Hung, J. K., & Treitz, P. (2020). Environmental land-cover classification for integrated watershed studies: Cape Bounty, Melville Island, Nunavut. *Arctic Science*, 6(4), 404–422. <http://dx.doi.org/10.1139/AS-2019-0029>
- Instituto da Conservação da Natureza e das Florestas. (2019). *Sub-regiões homogêneas dos Programas Regionais de Ordenamento Florestal (PROF)* [Homogeneous sub-regions of the Regional Forest Management Programs]. ICNF. https://geocatalogo.icnf.pt/metadados/prof_srh.html
- Instituto Nacional de Estatística. (2022). *Recenseamento agrícola* [Agricultural Census]. INE.

- Jansen, J. (2002). *Guia Geobotânico da Serra da Estrela* [Geobotanical Guide of Serra da Estrela]. Instituto da Conservação da Natureza.
- Jansen, J. (2011). *Managing Natura 2000 in a changing world: The case of the Serra da Estrela* (Portugal). [Doctoral dissertation, Radboud University Nijmegen]. Repository UBN. <https://repository.ubn.ru.nl/bitstream/handle/2066/90921/90921.pdf?sequence=1&isAllowed=y>
- Körner, C. (2004). Mountain biodiversity, its causes and function: An overview. *Ambio, Spec No 13*, 11–17.
- Machado, C., Cristina, M., & Sales, F. (2007). Expedições botânicas ao PNSE [Botanical expeditions to PNSE]. *Bloco de Notas do Centro de Interpretação da Serra da Estrela* [Bulletin of the Serra da Estrela Interpretation Centre], 20, 6–7.
- Meireles, C. (2010). *Flora e vegetação da Serra da Estrela aproximação fitossociológica da vertente meridional*. [Flora and vegetation of Serra da Estrela phytosociological approach of the southern slope]. [Dissertação de Doutoramento, Universidad de Jaén]. Dialnet. <https://dialnet.unirioja.es/servlet/tesis?codigo=153868>
- Meireles, C., & Pinto-Gomes, C. (2012). *Guia de campo do itinerário geobotânico à Serra da Estrela*. [Field guide for the geobotanical itinerary to Serra da Estrela]. Gobierno de la Extremadura. <http://hdl.handle.net/10174/7438>
- Meireles, C., Pinto-Gomes, C., Paiva-Ferreira, R., & Neiva, R. (2008). Valorização do Parque Natural da Serra da Estrela, desde o ponto de vista da flora e vegetação [Enhancement of the Serra da Estrela Natural Park, from the point of view of flora and vegetation]. In M. M. Redondo, M. T. Palácios, F. López & D. Sanches-Mata (Eds.), *Avances en Biogeografía* [Advances in Biogeography] (pp. 105–108). Universidad Autónoma de Madrid.
- Monteiro-Henriques, T., Martins, M., Cerdeira, J., Silva, P., Arsénio, P., Silva, Á., Bellu, A., & Costa, J. (2016). Bioclimatological mapping tackling uncertainty propagation: Application to mainland Portugal. *International Journal of Climatology*, 36, 400–411. <https://doi.org/10.1002/joc.4357>
- Mora, C. (2006). *Climas da Serra da Estrela: características regionais e particularidades locais dos planaltos e do alto do Vale do Zêzere* [Climates of Serra da Estrela: Regional characteristics and local particularities of the plateaus and the upper Zêzere Valley]. [Doctoral dissertation, Faculdade de Letras da Universidade de Lisboa]. Repositório da Universidade de Lisboa. <http://hdl.handle.net/10400.5/98023>
- Pauli, H., Gottfried, M., Dullinger, S., Abdaladze, O., Akhalkatsi, M., Benito Alonso, J. L., & Coldea, G. (2012). Recent plant diversity changes on Europe's mountain summits. *Science*, 336(6079), 353–355. <https://doi.org/10.1126/science.1219033>
- Pauli, H., Gottfried, M., & Grabherr, G. (2003). Effects of climate change on the alpine and nival vegetation of the Alps. *Journal of Mountain Ecology*, 7, 9–12. <https://www.cipra.org/en/publications/4846>
- Pinto da Silva, A. R., & Teles, A. N. (1980). *A Flora e a vegetação da Serra da Estrela* [Flora and vegetation of Serra da Estrela]. Serviço Nacional de Parques, Reservas e Património Paisagístico.
- Rapinel, S., Clément, B., Magnanon, S., Sellin, V., & Hubert-Moy, L. (2014). Identification and mapping of natural vegetation on a coastal site using a Worldview-2 satellite image. *Journal of Environmental Management*, 144, 236–246. <https://doi.org/10.1016/j.jenvman.2014.05.027>
- Rudley, D., DeSoto, L., Rodríguez-Echeverría, S., & Nabais, C. (2023). Climate effect on the growth and hydraulic traits of two shrubs from the top of a Mediterranean mountain. *Science of the Total Environment*, 902, 165911. <https://doi.org/10.1016/j.scitotenv.2023.165911>
- Santos, R. N., Rodrigues, T., Naughton, F., Schefuß, E., Oliveira, D., Moreno, J., ... & Hernández, A. (2024). Understanding the Atlantic influence on climate and vegetation dynamics in western Iberia over the last 2000 years. *Quaternary Science Reviews*, 337, 108796. <https://doi.org/10.1016/j.quascirev.2024.108796>
- Sanz-Elorza, M., Dana, E. D., González, A., & Sobrino, E. (2003). Changes in the high-mountain vegetation of the Central Iberian Peninsula as a probable sign of global warming. *Annals of Botany*, 92(2), 273–280. <https://doi.org/10.1093/aob/mcg130>
- Soncco, C. J. M. (2020). *Cartografia de ultra-alta resolução para monitorio y gestión de geossítios – Estrela Geopark* [Ultra-high resolution mapping for geosite monitoring and management – Estrela Geopark] [Doctoral dissertation, Universidade de Lisboa]. Repositório da Universidade de Lisboa. <http://hdl.handle.net/10451/44390>
- Tasser, E., Walde, J., Tappeiner, U., Teutsch, A., & Noggler, W. (2005). Land-use changes and natural reforestation in the Eastern Central Alps. *Agriculture, Ecosystems & Environment*, 113(1–4), 65–74. <http://dx.doi.org/10.1016/j.agee.2006.05.004>
- Van der Knaap, W. O., & van Leeuwen, J. (1994). Holocene vegetation, human impact, and climatic change in the Serra da Estrela, Portugal. Review of Palaeobotany and Palynology, 89(3–4), 153–221. [https://doi.org/10.1016/0034-6667\(95\)00048-0](https://doi.org/10.1016/0034-6667(95)00048-0)
- Vieira, G. (2004). *Geomorfologia dos planaltos e altos vales da Serra da Estrela: Ambientes frios do Plistocénico Superior e dinâmica atual* [Geomorphology of the plateaus and high valleys of Serra da Estrela: Cold environments of the

- Upper Pleistocene and current dynamics]. [Doctoral dissertation, Universidade de Lisboa]. Repositório da Universidade de Lisboa. <https://hdl.handle.net/10451/57057>
- Vieira, G., & Mora, C. (1998). General characteristics of the climate of the Serra da Estrela. In *Glacial and periglacial geomorphology of the Serra da Estrela: Guidebook for the fieldtrip, IGU Commission on Climate Change and Periglacial Environments* (pp. 26–36). CEG, University of Lisbon.
- Viviroli, D., Dürr, H. H., Messerli, B., Meybeck, M., & Weingartner, R. (2007). Mountains of the world, water towers for humanity: Typology, mapping, and global significance. *Water Resources Research*, 43(7), W07447. <https://doi.org/10.1029/2006WR005653>
- Wang, K., Zhang, L., Qiu, Y., Ji, L., Tian, F., Wang, C., & Wang, Z. (2015). Snow effects on alpine vegetation in the Qinghai-Tibetan Plateau. *International Journal of Digital Earth*, 8(1), 58-75. <https://doi.org/10.1080/17538947.2013.848946>
- Xie, Y., Sha, Z., & Yu, M. (2008). Remote sensing imagery in vegetation mapping: A review. *Journal of Plant Ecology*, 1(1), 9-23. <https://doi.org/10.1093/jpe/rtm005>

LEGAL DOCUMENTS

- Portugal. (1976). *Decreto-Lei n.º 557/76, de 16 de julho*. Diário da República, 1.ª série, n.º 165, 1562–1565. <https://diariodarepublica.pt/dr/detalhe/decreto-lei/557-1976-430620>